

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5 : C07D 215/44, 215/26, A61K 31/47		A1	(11) International Publication Number: WO 94/29274
		(43) International Publication Date: 22 December 1994 (22.12.94)	
(21) International Application Number: PCT/SE94/00552		(81) Designated States: AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LU, LV, MD, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 8 June 1994 (08.06.94)			
(30) Priority Data: 9302005-5 11 June 1993 (11.06.93) SE 9303970-9 30 November 1993 (30.11.93) SE			
(71) Applicant (for all designated States except US): AKTIEBO- LAGET ASTRA [SE/SE]; S-151 85 Södertälje (SE).		Published With international search report.	
(72) Inventor; and (75) Inventor/Applicant (for US only): STARKE, Carl, Ingemar [SE/SE]; Frigångsgatan 10, S-413 01 Göteborg (SE).			
(74) Agent: HÖGLUND, Lars; AB Astra, Patent Dept., S-151 85 Södertälje (SE).			

(54) Title: NEW ACTIVE COMPOUNDS

(57) Abstract

The present invention relates to novel quinoline derivatives of formula (I) which inhibit exogenously or endogenously stimulated gastric acid secretion and thus can be used in the prevention and treatment of gastrointestinal inflammatory diseases.



(I)

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

-1-

NEW ACTIVE COMPOUNDS

TECHNICAL FIELD

5 The present invention relates to novel compounds, and therapeutically acceptable salts thereof, which inhibit exogenously or endogenously stimulated gastric acid secretion and thus can be used in the prevention and treatment of gastrointestinal inflammatory diseases. In further aspects, the invention relates to compounds of the invention for use in therapy; to
10 processes for preparation of such new compounds; to pharmaceutical compositions containing at least one compound of the invention, or a therapeutically acceptable salt thereof, as active ingredient; and to the use of the active compounds in the manufacture of medicaments for the medical use indicated above.

BACKGROUND ART


20 Substituted quinoline derivatives that inhibit gastric acid secretion are known in the art, for example from EP-A1-259,174 and EP-A1-330,485.

DISCLOSURE OF THE INVENTION

25 It has surprisingly been found that compounds of the formula I, which are
4-amino-3-acylquinoline derivatives in which the quinoline is substituted in
the 8-position by alkylthioethoxy, alkylthiopropoxy, alkylsulfinylethoxy,
alkylsulfinylpropoxy, alkylsulfonylethoxy or alkylsulfonylpropoxy, are
effective as inhibitors of gastric acid secretion, and that they exert this
30 effect by inhibiting the gastrointestinal H^+, K^+ -ATPase.

In one aspect, the invention thus relates to compounds of the general formula I:

-2-



or a pharmaceutically acceptable salt thereof, wherein

15 R^1 is

(a) C₁-C₆ alkyl,
(b) C₃-C₆ cykloalkyl, or
(c) C₃-C₆, C₁-C₆ cykloalkylalkyl;

 R^2 is

20 (a) H,
(b) C₁-C₆ alkyl,
(c) C₁-C₆ alkoxy, or
(d) halogen;

R³ is C₁-C₆ alkyl;

25 \mathbb{R}^4 is

(a) H,
(b) C₁-C₄ alkyl,
(c) halogen, or
(d) OH;

30 m is an integer 2 or 3; and
 n is an integer 0, 1 or 2.

WO 94/29274

PCT/SE94/00552

—3—

Throughout the specification and the appended claims, a given chemical formula or name shall encompass all stereo and optical isomers and racemates thereof where such isomers exist, as well as pharmaceutically acceptable acid addition salts thereof and solvates thereof such as for instance hydrates.

The following definitions shall apply throughout the specification and the appended claims.

10 Unless otherwise stated or indicated, the term "C₁-C₆ alkyl" denotes a straight or branched alkyl group having from 1 to 6 carbon atoms. Examples of said lower alkyl include methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, t-butyl and straight- and branched-chain pentyl and hexyl.

15 Unless otherwise stated or indicated, the term "cycloalkyl" denotes a cyclic alkyl group having a ring size from C₃ to C₆, optionally additionally substituted by lower alkyl. Examples of said cycloalkyl include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, methylcyclohexyl and cycloheptyl.

20 Unless otherwise stated or indicated, the term "C₁-C₆ alkoxy" denotes a straight or branched alkoxy group having from 1 to 6 carbon atoms. Examples of said lower alkoxy include methoxy, ethoxy, n-propoxy, iso-propoxy, n-butoxy, iso-butoxy, sec-butoxy, t-butoxy and straight- and branched-chain pentoxy and hexoxy.

25 Unless otherwise stated or indicated, the term "halogen" shall mean fluorine, chlorine, bromine or iodine.

30 Depending on the process conditions and the starting materials, the end products of the formula I are obtained either in neutral or salt form. Both

WO 94/29274

PCT/SE94/00552

—4—

the free base and the salts of these end products are within the scope of the invention.

5 Acid addition salts of the new compounds may in a manner known *per se* be transformed into the free base using basic agents such as alkali or by ion exchange. The free base obtained may also form salts with organic or inorganic acids.

10 In the preparation of acid addition salts, preferably such acids are used which form suitably therapeutically acceptable salts. Examples of such acids are hydrohalogen acids, sulfonic acids, phosphoric acid, nitric acid, aliphatic, alicyclic, aromatic or heterocyclic carboxyl or sulfonic acids, such as formic acid, acetic acid, propionic acid, succinic acid, glycolic acid, lactic acid, malic acid, tartaric acid, citric acid, ascorbic acid, maleic acid,
15 hydroxymaleic acid, pyruvic acid, p-hydroxybenzoic acid, embonic acid, methanesulfonic acid, ethanesulfonic acid, hydroxyethanesulfonic acid, halogenbensenesulfonic acid, toluenesulfonic acid or naphthalenesulfonic acid.

20 Preferred compounds of the formula I are those wherein
R¹ is CH₃, CH₂CH₃, CH₂CH₂CH₃, CH(CH₃)₂, cyclopropyl or cyclopropylmethyl;
R² is CH₃, CH₂CH₃, CH₂CH₂CH₃, CH(CH₃)₂, CH(CH₃)CH₂CH₃, OCH₃, OCH₂CH₃, or halogen;
25 R³ is CH₃, CH₂CH₃, CH(CH₃)₂ or CH₂CH₂CH₃; and
R⁴ is H, CH₃, CH₂CH₃, halogen or OH.

More preferred compounds of the formula I are those wherein
R¹ is CH₂CH₃ or CH₂CH₂CH₃;
30 R² is CH₃, CH₂CH₃, CH(CH₃)₂, OCH₃, or Cl;
R³ is CH₃, CH₂CH₃ or CH₂CH₂CH₃; and
R⁴ is H, CH₃, F, Cl or OH.

WO 94/29274

PCT/SE94/00552

—5—

The most preferred compound of the invention is that wherein R^1 is $CH_2CH_2CH_3$; R^2 and R^3 is CH_3 ; R^4 is H; m is 2; and n is 1; i.e. the compound 3-butyryl-4-(2-methylphenylamino)-8-(2-methylsulfinylethoxy)quinoline.

5


In a further aspect, the invention relates to compounds of the formula I for use in therapy.

Preparation

10 The present invention also provides processes for the manufacture of the compounds with the general formula I. Such compounds are prepared in the following way:

(A) A compound of the general formula II

15




II

20

wherein R^2 and R^4 are as defined above is reacted with a compound of the general formula III

25



III

30

WO 94/29274

PCT/SE94/00552

—6—

wherein R^1 , R^3 , m and n are as defined above and X is a leaving group, such as a halide, tosyloxy or mesyloxy.

5 Compounds of formula III are novel and represent as such a further aspect of the invention.

This reaction is conducted with or without a solvent. When a solvent is used, it is preferably a solvent such as acetonitrile, tetrahydrofuran, toluene or dimethyl formamide.

10

When the reaction is conducted with a solvent, the reaction temperature usually ranges from about 20°C to about the boiling point of the solvent used, more preferably from about 20°C to about 110°C. The reaction time usually ranges from about 1 hour to about 24 hours.

15

When the reaction is conducted without a solvent, the reaction temperature usually ranges from about 30°C to about 170°C. The reaction time usually ranges from 15 minutes to about 2 hours.

20

(B) Compounds of the formula I wherein R^1 , R^2 , R^3 , R^4 and m are as defined above under (A) and n is 1 or 2 can be prepared by oxidizing a compound of formula I wherein R^1 , R^2 , R^3 , R^4 and m are as defined above under (A) and n is 0.

25

This oxidation may be carried out by using an oxidizing agent such as sodium hypochlorite, nitric acid, hydrogen peroxide, (optionally in presence of vanadium compounds), peracids, peresters, ozone, dinitrogen tetraoxide, iodosobenzene, N-halosuccinimide, 1-chlorobenzotriazole, t-butylhypochlorite, diazabicyclo-[2,2,2]-octane

30

bromine complex, sodium metaperiodate, selenium dioxide, manganese dioxide, chromic acid, ceric ammonium nitrate, bromine, chlorine, and sulfonyl chloride. The oxidation takes place in a solvent such as halogenated hydrocarbons, alcohols, ethers or ketones.

WO 94/29274

PCT/SE94/00552

—7—

The oxidation may also be carried out enzymatically by using an oxidizing enzyme or microbiotically by using a suitable microorganism.

5 Compounds of the general formula II are commercially available or can be prepared by known methods.

Compounds of the general formula III can be prepared according to known methods and according to the Examples below.

10 Use

In a further aspect, the invention relates to the use of a compound as defined above for the manufacture of a medicament for the inhibition of gastric acid secretion, or for the treatment of gastrointestinal inflammatory diseases.

15

In a more general sense, the compounds of the invention may be used for prevention and treatment of gastrointestinal inflammatory diseases, and gastric acid-related diseases in mammals including man, such as gastritis, gastric ulcer, duodenal ulcer, reflux esophagitis and Zollinger-Ellison syndrom.

20

Furthermore, the compounds may be used for treatment of other gastrointestinal disorders where gastric antisecretory effect is desirable, e.g. in patients with gastrinomas, and in patients with acute upper gastrointestinal bleeding. They may also be used in patients in intensive care situations, and pre-and postoperatively to prevent acid aspiration and stress ulceration.

25

Pharmaceutical formulations

30 In yet a further aspect, the invention relates to pharmaceutical compositions containing at least one compound of the invention, or a therapeutically acceptable salt thereof, as active ingredient.

WO 94/29274

PCT/SE94/00552

-8-

The compounds of the invention can also be used in formulations together with other active ingredients, e.g. for the treatment or prophylaxis of conditions involving infection by *Helicobacter pylori* of human gastric mucosa. Such other active ingredients may be antimicrobial agents,

5 especially:

β -lactam antibiotics such as amoxicillin, ampicillin, cephalothin, cefaclor or cefixime; or

macrolides such as erythromycin, or clarithromycin; or

tetracyclines such as tetracycline or doxycycline; or

10 aminoglycosides such as gentamycin, kanamycin or amikacin; or

quinolones such as norfloxacin, ciprofloxacin or enoxacin; or

others such as metronidazole, nitrofurantoin or chloramphenicol;

or preparations containing bismuth salts such as bismuth subcitrate, bismuth subsalicylate, bismuth subcarbonate, bismuth subnitrate or

15 bismuth subgallate.

For clinical use, the compounds of the invention are formulated into pharmaceutical formulations for oral, rectal, parenteral or other mode of administration. The pharmaceutical formulation contains a compound of

20 the invention in combination with one or more pharmaceutically acceptable ingredients. The carrier may be in the form of a solid, semi-solid or liquid diluent, or a capsule. These pharmaceutical preparations are a further object of the invention. Usually the amount of active compounds is between 0.1-95% by weight of the preparation, preferably between 0.2-20%

25 by weight in preparations for parenteral use and preferably between 1 and 50% by weight in preparations for oral administration.

In the preparation of pharmaceutical formulations containing a compound of the present invention in the form of dosage units for oral administration

30 the compound selected may be mixed with solid, powdered ingredients, such as lactose, saccharose, sorbitol, mannitol, starch, amylopectin, cellulose derivatives, gelatin, or another suitable ingredient, as well as with disintegrating agents and lubricating agents such as magnesium stearate,

WO 94/29274

PCT/SE94/00552

-9-

calcium stearate, sodium steryl fumarate and polyethylene glycol waxes.

The mixture is then processed into granules or pressed into tablets.

Soft gelatine capsules may be prepared with capsules containing a mixture of the active compound or compounds of the invention, vegetable oil, fat, or other suitable vehicle for soft gelatine capsules. Hard gelatine capsules may contain granules of the active compound. Hard gelatine capsules may also contain the active compound in combination with solid powdered ingredients such as lactose, saccharose, sorbitol, mannitol, potato starch, corn starch, amylopectin, cellulose derivatives or gelatine.

10

Dosage units for rectal administration may be prepared (i) in the form of suppositories which contain the active substance mixed with a neutral fat base; (ii) in the form of a gelatine rectal capsule which contains the active substance in a mixture with a vegetable oil, paraffin oil or other suitable vehicle for gelatine rectal capsules; (iii) in the form of a ready-made micro enema; or (iv) in the form of a dry micro enema formulation to be reconstituted in a suitable solvent just prior to administration.

15

Liquid preparations for oral administration may be prepared in the form of syrups or suspensions, e.g. solutions or suspensions containing from 0.2% to 20% by weight of the active ingredient and the remainder consisting of sugar or sugar alcohols and a mixture of ethanol, water, glycerol, propylene glycol and polyethylene glycol. If desired, such liquid preparations may contain colouring agents, flavouring agents, saccharine and carboxymethyl cellulose or other thickening agent. Liquid preparations for oral administration may also be prepared in the form of a dry powder to be reconstituted with a suitable solvent prior to use.

20

25

Solutions for parenteral administration may be prepared as a solution of a compound of the invention in a pharmaceutically acceptable solvent, preferably in a concentration from 0.1% to 10% by weight. These solutions may also contain stabilizing ingredients and/or buffering ingredients and are dispensed into unit doses in the form of ampoules or vials. Solutions

30

-10-

for parenteral administration may also be prepared as a dry preparation to be reconstituted with a suitable solvent extemporaneously before use.

5 The typical daily dose of the active substance varies within a wide range and will depend on various factors such as for example the individual requirement of each patient, the route of administration and the disease. In general, oral and parenteral dosages will be in the range of 5 to 1000 mg per day of active substance.

10

EXAMPLES

1. PREPARATION OF COMPOUNDS OF THE INVENTION

15 Example 1

Preparation of 3-butyryl-4-(2-methylphenylamino)-8-(2-methylthioethoxy)quinoline

20 A mixture of 3-butyryl-4-chloro-8-(2-methylthioethoxy)quinoline (0.67 g, 2.1 mmol) and o-toluidine (0.24 g, 2.3 mmol) in acetonitrile was heated to 55°C and stirred for 3.5 h. The solvent was evaporated and the residue was partitioned between methylene chloride and a saturated sodium bicarbonate solution. The organic layer was dried over sodium sulfate and evaporated. The residue was triturated with diisopropylether. The
25 precipitated product was filtered off and washed with diisopropyl ether giving 0.55 g (66%) of the title compound.

30 (¹H-NMR, 500 MHz, CDCl₃) 1.05 (t,3H), 1.84 (m,2H), 2.25 (s,3H), 2.35 (s,3H), 3.10 (m,4H), 4.34 (t,2H), 6.89 (d,1H), 6.95-7.15 (m,5H), 7.27 (d,1H), 9.26 (s,1H), 11.84 (s,1H).

WO 94/29274

PCT/SE94/00552

-11-

Example 2Preparation of 3-butyryl-4-(2-methylphenylamino)-8-(2-methylsulfinylethoxy)quinoline

5 3-Butyryl-4-(2-methylphenylamino)-8-(2-methylthioethoxy)quinoline (0.15 g, 0.38 mmol) was dissolved in methylene chloride (3 ml) and cooled to -20°C. A solution of 71% m-CPBA (0.089 g, 0.36 mmol) in 1 ml of methylene chloride was added dropwise. The temperature was allowed to rise to room temperature whereafter the solution was stirred for 15 min at room temperature. The reaction mixture was washed with a saturated sodium bicarbonate solution. The organic layer was dried over sodium sulfate and evaporated. Chromatography with methylene chloride:methanol 10:1 as the eluent gave 0.064 g (41%) of the desired product.

15 ¹H-NMR, 300 MHz, CDCl₃) 1.04 (t,3H), 1.82 (m,2H), 2.34 (s,3H), 2.80 (s,3H), 3.08 (t,2H), 3.21 (m,1H), 3.44 (m,1H), 4.62 (m,2H), 6.89 (d,1H), 6.94-7.16 (m,5H), 7.28 (d,1H), 9.20 (s,1H), 11.82 (s,1H).

Example 3Preparation of 3-butyryl-4-(2-methylphenylamino)-8-(2-methylsulfonylethoxy)quinoline

25 3-Butyryl-4-(2-methylphenylamino)-8-(2-methylthioethoxy)quinoline (0.037 g, 0.092 mmol) was dissolved in methylene chloride (1.5 ml) and cooled to -20°C. A solution of 71% m-CPBA (0.047 g, 0.19 mmol) in 0.5 ml methylene chloride was added dropwise. The temperature was allowed to rise to room temperature whereafter the solution was stirred for 30 min at room temperature and the organic layer was washed with a saturated sodium bicarbonate solution. The organic layer was dried over sodium sulfate and evaporated. Trituration with diisopropyl ether and the product crystallized. After chromatography of the precipitate with methylene

30

WO 94/29274

PCT/SE94/00552

—12—

chloride:ethyl acetate 1:1 as the eluent, at the end pure ethyl acetate, 0.022 g (56%) of the title compound was isolated.

5 $^1\text{H-NMR}$, 500 MHz, CDCl_3) 1.07 (t,3H), 1.84 (m,2H), 2.35 (s,3H), 3.10 (t,2H), 3.39 (s,3H), 3.62 (t,2H), 4.61 (t,2H), 6.89 (d,1H), 6.94-7.17 (m,5H), 7.28 (m,1H), 9.15 (s,1H), 11.86 (s,1H).

Example 4

10 Preparation of 3-butyryl-4-(2-isopropylphenylamino)-8-(2-methylthioethoxy)quinoline

A mixture of 3-butyryl-4-chloro-8-(2-methylthioethoxy)quinoline (400 mg, 1.23 mmol) and 2-isopropylaniline (1.0 g, 7.4 mmol) was heated to 150°C for 30 min. The mixture was diluted with CHCl_3 and extracted with 2 N HCl. The organic phase was washed with a saturated sodium bicarbonate solution. The organic layer was dried over Na_2SO_4 and evaporated. The residue was chromatographed (SiO_2 ; CH_2Cl_2 :MeOH 95:5) yielding 340 mg (80.5%) of the desired product.

20 $^1\text{H-NMR}$, 300 MHz, CDCl_3): 1.0 (t,3H), 1.1 (d,3H), 1.2 (d,3H), 1.75 (m,2H), 2.15 (s,3H), 3.1 (m,3H), 3.2 (t,2H), 4.3 (t,2H), 6.8 (d,1H), 7.0-7.2 (m,4H), 7.4 (m,2H), 9.4 (d,1H).

Example 5

25 Preparation of 3-butyryl-4-(2-isopropylphenylamino)-8-(2-methylsulfinylethoxy)quinoline.

3-Butyryl-4-(2-isopropylphenylamino)-8-(2-methylthioethoxy)quinoline (0.22 g, 0.52 mmol) was dissolved in methylene chloride (15 ml) and added to a mixture of 0.093g NaHCO_3 in 15 ml H_2O . A solution of 71% m-CPBA (0.12 g, 0.50 mmol) in 7 ml methylene chloride was added dropwise at 4°C. The solution was stirred for one hour at this temperature. The reaction mixture was washed with a saturated sodium bicarbonate solution. The organic

-13-

layer was dried over Na_2SO_4 and evaporated. Chromatography with methylene chloride:methanol 10:1 as eluent gave 0.130 g (57%) of the desired product.

- 5 $^1\text{H-NMR}$, 300 MHz, CDCl_3). 1.0 (t,3H), 1.3 (m,6H), 1.8 (m,2H), 2.78 (s,3H), 3.08 (t,2H), 3.2 (m,6H), 3.35 (m,1H), 3.45 (m,1H), 4.6 (m,2H), 6.82 (d,1H), 6.86 (t,1H), 7.05 (m,3H), 7.2 (t,1H), 7.38 (d,1H), 9.18 (s,1H), 11.8 (s,1H).

Example 6

- 10 Preparation of 3-butyryl-4-(2-isopropylphenylamino)-8-(2-methylsulfonylethoxy)quinoline.

A mixture of 3-butyryl-4-(2-isopropylphenylamino)-8-(2-methylthioethoxy)quinoline (0.22 g, 0.52 mmol) in 15 ml methylene
15 chloride and NaHCO_3 (0.186 g, 2.2 mmol) in 15 ml H_2O was cooled to 4°C . A solution of 70% m-CPBA (0.24 g, 1.0 mmol) in 7 ml methylene chloride was added dropwise. After stirring for 1 h at 4°C , the organic layer was dried over Na_2SO_4 and evaporated. Chromatography (SiO_2 ; CH_2Cl_2 :MeOH 90:10) gave 16 mg (6.8%) of the desired product.

- 20 $^1\text{H-NMR}$, 300 MHz, CDCl_3): 1.05 (t,3H), 1.2 (m,6H), 1.8 (m,2H), 3.1 (t,2H), 3.35 (s,3H), 3.45 (m,1H), 3.65 (m,2H), 4.6 (m,2H), 6.85 (d,2H), 6.9-7.1 (m,4H), 7.4 (d,1H), 9.1(s,1H), 11.8 (s,1H).

25 Example 7

Preparation of 3-propanoyl-4-(2-methylphenylamino)-8-(2-methylthioethoxy)quinoline.

- 30 A mixture of 3-propanoyl-4-chloro-8-(2-methylthioethoxy)quinoline (0.60 g, 1.9 mmol) and o-toluidine (0.25 g, 2.3 mmol) in acetonitrile was heated to 55°C and stirred for 3.5 h. The solvent was evaporated and the residue was partitioned between methylene chloride and a 10% sodium carbonate solution. The organic layer was dried over sodium sulfate and evaporated.

-14-

Chromatography (SiO₂; CH₂Cl₂:EtOAc 60:40) gave 0.45 g (61%) of the title compound.

(¹H-NMR, 300 MHz, CDCl₃) 1.26 (t, 3H), 2.21 (s, 3H), 2.33 (s, 3H), 3.06 (t, 2H),
 5 3.10 (q, 2H), 4.33 (t, 2H), 6.87 (d, 1H), 6.96-7.12 (m, 5H), 7.25 (d, 1H), 9.26
 (s, 1H), 11.78 (s, 1H).

Example 8

10 Preparation of 3-propanoyl-4-(2-methylphenylamino)-8-(2-methylsulfinylethoxy)quinoline.

3-Propanoyl-4-(2-methylphenylamino)-8-(2-methylthioethoxy)quinoline
 (0.10 g, 0.26 mmol) was dissolved in methylene chloride (5 ml). NaHCO₃
 (45 mg) in H₂O (5 ml) was added. The mixture was cooled to 4°C. A
 15 solution of 71% m-CPBA (0.062 g, 0.25 mmol) in methylene chloride (5 ml)
 was added dropwise. After stirring for 1 h at 2-4°C, the organic layer was
 washed with a saturated sodium bicarbonate solution. The organic layer
 was dried over sodium sulfate and evaporated. Chromatography (SiO₂;
 CH₂Cl₂:EtOH 90:10) gave 50 mg (48%) of the desired product.

20 (¹H-NMR, 300 MHz, CDCl₃) 1.26 (t, 3H), 2.33 (s, 3H), 2.78 (s, 3H), 3.10-3.50
 (m, 4H), 4.61 (m, 2H), 6.85 (d, 1H), 6.92-7.11 (m, 5H), 7.28 (d, 1H), 9.18 (s, 1H),
 11.81 (s, 1H).

25 Example 9

Preparation of 3-propanoyl-4-(2-methylphenylamino)-8-(2-methylsulfonylethoxy)quinoline.

3-Propanoyl-4-(2-methylphenylamino)-8-(2-methylthioethoxy)quinoline
 30 (0.12 g, 0.32 mmol) was dissolved in methylene chloride (5 ml). NaHCO₃
 (110 mg) in H₂O (10 ml) was added. The mixture was cooled to 4°C. A
 solution of 71% m-CPBA (0.17 g, 0.69 mmol) in methylene chloride (5 ml)
 was added dropwise. After stirring for 1 h at 2-4°C, the organic layer was

-15-

washed with a saturated sodium bicarbonate solution. The organic layer was dried over sodium sulfate and evaporated. Chromatography (SiO₂; CH₂Cl₂:EtOAc 50:50) gave 0.060 g (45%) of the title compound.

5 ¹H-NMR, 300 MHz, CDCl₃) 1.26 (t,3H), 2.34 (d,3H), 3.15 (q,2H), 3.37 (s,3H), 3.61 (t,2H), 4.58 (t,2H), 6.86 (d,1H), 6.95-7.11 (m,5H), 7.26 (d,1H), 9.13 (s,1H), 11.81 (s,1H).

Example 10

10 Preparation of 3-propanoyl-4-(2-ethylphenylamino)-8-(2-methylthioethoxy)quinoline

A mixture of 3-propanoyl-4-chloro-8-(2-methylthioethoxy)quinoline (0.093 g, 0.34 mmol) and 2-ethyl aniline (0.048 g, 0.39 mmol) in acetonitrile (1 ml) was heated to 65°C and stirred 4.0 h. The solvent was evaporated and the residue was partitioned between methylene chloride and a saturated sodium bicarbonate solution. The organic layer was dried over sodium sulfate and evaporated. The residue was chromatographed (SiO₂; ethyl acetate) yielding 40 mg (30%) of the desired product.

20 ¹H-NMR, 300 MHz, CDCl₃) 1.22-1.30 (m,6H), 2.22 (s,3H), 2.76 (q,2H) 3.06 (t,2H), 3.15 (q,2H), 4.34 (t,2H), 6.82 (d,1H), 6.91-7.06 (m,2H), 7.14 (m,1H), 7.28 (m,1H), 9.21 (s,1H), 11.83 (s,1H).

25 Examples 11 and 12

Preparation of 3-propanoyl-4-(2-ethylphenylamino)-8-(2-methylsulfinylethoxy)quinoline (Example 11) and 3-propanoyl-4-(2-ethylphenylamino)-8-(2-methylsulfonylethoxy)quinoline (Example 12)

30 3-Propanoyl-4-(2-ethylphenylamino)-8-(2-methylthioethoxy)quinoline (0.022 g, 0.056 mmol) was dissolved in methylene chloride (0.7 ml). NaHCO₃ (12 mg) in H₂O (0.7 ml) was added. The mixture was cooled to 4°C. A solution of 71% m-CPBA (0.017 g, 0.07 mmol) in methylene chloride (0.5

WO 94/29274

PCT/SE94/00552

-16-

ml) was added dropwise. After stirring for 1 h at 2-4°C, the organic layer was washed with a saturated sodium bicarbonate solution. The organic layer was dried over sodium sulfate and evaporated. Chromatography (SiO₂; CH₂:MeOH 90:10) gave 8 mg (35%) of the compound according to Example 11 and 10 mg (42%) of the compound according to Example 12.

Example 11: (¹H-NMR, 300 MHz, CDCl₃): 1.25 (m, 6H), 2.73-2.81 (m, 5H), 3.15 (q, 2H), 3.22 (m, 1H), 3.41-3.49 (m, 1H), 4.62 (m, 2H), 6.84 (d, 1H), 6.93-7.19 (m, 5H), 7.31 (d, 1H), 9.19 (s, 1H), 11.88 (s, 1H).

Example 12: (¹H-NMR, 300 MHz, CDCl₃): 1.29 (m, 6H), 2.77 (q, 2H), 3.16 (q, 2H), 3.37 (s, 3H), 3.61 (t, 2H), 4.60 (t, 2H), 6.85 (d, 1H), 6.94-7.20 (m, 5H), 7.31 (d, 1H), 9.14 (s, 1H), 11.90 (s, 1H).

Example 13

Preparation of 3-butyryl-4-(4-fluoro-2-methylphenylamino)-8-(2-methylthioethoxy)quinoline

A mixture of 3-butyryl-4-chloro-8-(2-methylthioethoxy)quinoline (2.75 g, 8.2 mmol) and 4-fluoro-2-methylaniline (1.34 g, 10.7 mmol) in acetonitrile (20 ml) was refluxed for 8 h. The solution was cooled and 1.52 g crystallized product was filtered off. The filtrate was evaporated and chromatographed (SiO₂; CH₂Cl₂:MeOH 95:5) yielding 0.4 g of the desired product. Total yield: 1.92 g (57%).

(¹H-NMR, 300 MHz, CDCl₃) 1.05 (t, 3H), 1.85 (m, 2 H), 2.20 (s, 3 H), 2.30 (s, 3H), 3.05 (m, 4H), 4.35 (t, 2H), 6.75-6.90 (m, 2H), 7.00 (m, 4H), 9.20 (s, 1H), 11.80 (s, 1H).

Examples 14 and 15

Preparation of 3-butyryl-4-(4-fluoro-2-methylphenylamino)-8-(2-methylsulfinylethoxy)quinoline (Example 14) and 3-butyryl-4-(4-fluoro-2-methylphenylamino)-8-(2-methylsulfonylethoxy)quinoline (Example 15)

WO 94/29274

PCT/SE94/00552

-17-

3-butyryl-4-(4-fluoro-2-methylphenylamino)-8-(2-methylthioethoxy) quinoline (1.6 g, 3.88 mmol) was dissolved in methylene chloride (35 ml). 0.3 M NaHCO₃-solution (36 ml) was added. The mixture was cooled to 4°C. A solution of 70% m-CPBA (1.22 g, 5.04 mmol) in methylene chloride
5 (16 ml) was added dropwise. After stirring for 1 h at 2-4°C, the organic layer was washed with 0.3 M NaHCO₃-solution. The organic layer was dried over Na₂SO₄ and evaporated. Chromatography (SiO₂; CH₂Cl₂:MeOH 95:5) gave 1.55 g (93%) of the compound according to Example 14 and 0.31 g (18%) of the compound according to Example 15.

10

Example 14: (¹H-NMR, 300 MHz, CDCl₃) 0.95 (t, 3H), 1.75 (m, 2H), 2.20 (s, 3H), 2.70 (s, 3H), 3.00 (t, 2H), 3.10 (m, 1H), 3.30-3.40 (m, 1H), 4.50 (m, 2H), 6.70 (m, 1H), 6.75 (m, 1H), 6.85-6.95 (m, 3H), 7.00 (m, 1H), 9.10 (s, 1H), 11.85 (s, 1H).

15

Example 15: (¹H-NMR, 300 MHz, CDCl₃) : 1.05 (t, 3H), 1.80 (m, 2H), 2.30 (s, 3H), 3.10 (t, 2H), 3.40 (s, 3H), 3.60 (t, 2H), 4.60 (t, 2H), 6.75-6.80 (m, 1H), 6.85-6.90 (m, 1H), 6.95-7.05 (m, 4H), 9.15 (s, 1H), 11.85 (s, 1H).

20

Example 16

Preparation of 3-propanoyl-4(2-isopropylphenylamino)-8-(2-methylthioethoxy)quinoline

A mixture of 3-propanoyl-4-chloro-8-(2-methylthioethoxy)quinoline (305
25 mg, 1 mmol) and 2-isopropylaniline (1 ml) in 25 ml acetonitrile was refluxed overnight. The solvent was evaporated and the residue was partitioned between methylene chloride and a saturated sodium bicarbonate solution. The organic layer was dried over sodium sulfate and evaporated. The residue was chromatographed on a prep-TLC (methylene
30 chloride : ethyl acetate 1:1) yielding 30 mg (8%) of the desired product.

WO 94/29274

PCT/SE94/00552

-18-

(¹H-NMR, 300 MHz, CDCl₃) 1.3 (m, 9H), 2.25 (s, 3H), 3.05 (t, 2H), 3.15 (q, 2H), 3.4 (m, 1H), 4.3 (t, 2H), 6.8 (d, 1H), 7.0 (m, 4H), 7.2 (t, 1H), 7.4 (d, 1H), 9.2 (s, 1H).

5 Example 17

Preparation of 3-butyryl-4-(2-ethylphenylamino)-8-(2-methylthioethoxy)quinoline

10 A mixture of 3-butyryl-4-chloro-8-(2-methylthioethoxy)quinoline (3.69 g, 9.84 mmol) and 2-ethylaniline (1.55 g, 12.8 mmol) in acetonitrile (20 ml) was refluxed for 6 h. The solution was evaporated. Chromatography (SiO₂; CH₂Cl₂ : MeOH 97:3) gave 2.79 g (69%) of the desired product.

15 (¹H-NMR, 300 MHz, CDCl₃) 1.00 (t, 3H), 1.25 (t, 3H), 1.80 (m, 2H), 2.20 (s, 3H), 2.80 (m, 2H), 3.10 (m, 4H), 4.35 (t, 2H), 6.85 (m, 1H), 6.90-7.10 (m, 4H), 7.15 (t, 1H), 7.30 (m, 1H), 9.20 (s, 1H).

Examples 18 and 19

20 Preparation of 3-butyryl-4-(2-ethylphenylamino)-8-(2-methylsulfinylethoxy)-quinoline (Example 18) and 3-butyryl-4-(2-ethylphenylamino)-8-(2-methylsulfonylethoxy)quinoline (Example 19).

25 3-butyryl-4-(2-ethylphenylamino)-8-(2-methylthioethoxy)quinoline (1.98 g, 4.85 mmol) was dissolved in methylene chloride (40 ml). 0.3 M NaHCO₃ solution (45 ml) was added. The mixture was cooled to 4°C. A solution of 70% m-CPBA (1.54 g, 6.31 mmol) in methylene chloride (20 ml) was added dropwise. After stirring for 1 h at 2-4°C, the organic layer was washed with 0.3 M NaHCO₃ solution. The organic layer was dried over NaSO₄ and evaporated. Chromatography (SiO₂; CH₂Cl₂:MeOH 97:3) gave 0.62 g (30%)
30 of Example 18 and 0.39 g (18%) of Example 19.

Example 18: (¹H-NMR, 300 MHz, CDCl₃) : 1.05 (t, 3H), 1.30 (t, 3H), 1.85 (m, 2H), 2.75 (m, 2H), 2.85 (s, 3H), 3.05 (t, 2H), 3.20-3.30 (m, 1H), 3.40-3.50

WO 94/29274

PCT/SE94/00552

-19-

(m, 1H), 4.65 (m, 2H), 6.85 (m, 1H), 6.90-7.00 (m, 1H), 7.05-7.10 (m, 3H), 7.15 (t, 1H), 7.30 (m, 1H), 9.20 (s, 1H), 11.95 (s, 1H).

5 Example 19: (^1H -NMR, 300 MHz, CDCl_3) : 1.05 (t, 3H), 1.30 (t, 3H), 1.85 (m, 2H), 2.75 (m, 2H), 3.10 (t, 2H), 3.40 (s, 3H), 3.65 (m, 2H), 4.60 (m, 2H), 6.85 (d, 1H), 6.95-7.10 (m, 4H), 7.20 (t, 1H), 7.30 (m, 1H), 9.10 (s, 1H).

Example 20

10 Preparation of 3-propanoyl-4-(2-methylphenylamino)-8-(2-propylthioethoxy)quinoline.

3-Propanoyl-4-chloro-8-(2-propylthioethoxy)quinoline (2.5 g, 7.40 mmol) and o-toluidine (0.95 g, 8.86 mmol) was refluxed in acetonitrile (10 ml) for 2 h. The solvent was evaporated and the residue was partitioned between 15 methylene chloride and 10% Na_2CO_3 solution. The organic layer was dried over Na_2SO_4 and evaporated. The crude product was purified by column chromatography (methylene chloride : ethyl acetate 80:20). 1.8 g (60%) of the title compound was obtained.

20 (^1H -NMR, 300 MHz, CDCl_3) 1.00 (t, 3H), 1.28 (t, 3H) 1.66 (m, 2H), 2.35 (s, 3H) 2.62 (t, 2H), 3.09 (t, 2H), 3.14 (q, 2H), 4.32 (t, 2H), 6.80-7.25 (m, 7H), 9.22 (s, 1H), 11.76 (s, 1H).

Example 21

25 Preparation of 3-propanoyl-4-(2-methylphenylamino)-8-(2-propylsulfinylethoxy)quinoline.

3-Propanoyl-4-(2-methylphenylamino)-8-(2-propylthioethoxy)quinoline (0.5 g, 1.22 mmol) was dissolved in 10 ml methylene chloride. A solution 30 of sodium bicarbonate (250 mg, 3.0 mmol) in 10 ml water was added. The mixture was cooled to 2-4°C. A solution of 70% m-CPBA (295 mg, 1.20 mmol) in 10 ml methylene chloride was added dropwise during 10 min. The temperature was allowed to rise to room temperature and the mixture

WO 94/29274

PCT/SE94/00552

-20-

was stirred 30 min at this temperature. The organic layer was dried over Na_2SO_4 and evaporated. The residue was purified by column chromatography (methylene chloride : ethanol 90:10) to give 380 mg (73%) of the title compound.

5

(^1H -NMR, 300 MHz, CDCl_3) 1.11 (t, 3H), 1.28 (t, 3H), 1.87 (m, 2H), 2.35 (s, 3H), 2.89 (m, 2H), 3.10-3.20 (m, 3H), 3.40 (m, 1H), 4.63 (q, 2H), 6.90-7.40 (m, 7H), 9.19 (s, 1H), 11.85 (s, 1H).

10

Example 22

Preparation of 3-propanoyl-4-(2-methylphenylamino)-8-(2-propylsulfonyl-ethoxy)quinoline.

15

3-propanoyl-4-(2-methylphenylamino)-8-(2-propylthioethoxy)quinoline (500 mg, 1.22 mmol) was dissolved in 10 ml methylene chloride. A solution of sodium bicarbonate (500 mg, 5.95 mmol) in 10 ml water was added. The mixture was cooled to 2-4°C. A solution of 70% m-CPBA (600 mg, 2.43 mmol) in 10 ml methylene chloride was added dropwise. The temperature was allowed to rise to room temperature and the stirring was continued for 30 min at this temperature. The methylene chloride layer was separated and washed with water. The organic layer was then dried over Na_2SO_4 and evaporated. The crude product was purified by column chromatography (methylene chloride : ethylacetate 50:50). 340 mg (63 %) of the title compound was obtained.

25

(^1H -NMR, 300 MHz, CDCl_3) 1.13 (t, 3H), 1.27 (t, 3H), 1.96 (m, 2H), 2.35 (s, 3H), 3.16 (q, 2H), 3.47 (t, 2H), 3.58 (t, 2H), 4.58 (t, 2H), 6.85-7.25 (m, 7H), 9.12 (s, 1H), 11.81 (s, 1H).

30

Example 23

Preparation of 3-propanoyl-4-(2-methylphenylamino)-8-(3-propylthio-propoxy)quinoline.

WO 94/29274

PCT/SE94/00552

-21-

3-Propanoyl-4-chloro-8-(3-propylthiopropoxy)quinoline (2.0 g, 5.7 mmol) and o-toluidine (0.7g, 6.5 mmol) was refluxed in acetonitrile (10 ml) for 2 h. The solvent was evaporated and the residue was partitioned between methylene chloride and 10% Na₂CO₃ solution. The organic layer was dried
5 over Na₂SO₄ and evaporated. The residue was purified by column chromatography (methylene chloride : ethyl acetate 70:30). 1.3 g (54 %) of the title compound was obtained.

(¹H-NMR, 300 MHz, CDCl₃) 0.94 (t, 3H), 1.26 (t, 3H), 1.57 (m, 2H), 2.25 (m,
10 2H), 2.34 (s, 3H), 2.49 (t, 2H), 2.75 (t, 2H), 3.15 (q, 2H), 4.27 (t, 2H) 6.83-7.23 (m, 7H), 9.22 (s, 1H), 11.73 (s, 1H).

Example 24

Preparation of 3-propanoyl-4-(2-methylphenylamino)-8-(3-propylsulfinyl-
15 propoxy)quinoline.

3-Propanoyl-4-(2-methylphenylamino)-8-(3-propylthiopropoxy)quinoline (200 mg, 0.47 mmol) was dissolved in 5 ml methylene chloride. A solution of sodium bicarbonate (80 mg, 0.95 mmol) in 5 ml water was added. The
20 mixture was cooled to 2-4°C. A solution of 70% m-CPBA (115mg, 0.47 mmol) in 5 ml methylene chloride was added dropwise during 10 min. The temperature was allowed to rise and the mixture was stirred for 30 min at room temperature. The methylene chloride layer was separated and washed with water. The organic layer was then dried over Na₂SO₄ and
25 evaporated. The crude product was purified by column chromatography (methylene chloride : ethanol 95:5). 160 mg (77%) of the title compound was obtained.

(¹H-NMR, 300 MHz, CDCl₃) 1.06 (t, 3H), 1.27 (t, 3H), 1.79 (m, 2H), 2.33
30 (s, 3H), 2.49 (m, 2H), 2.60-2.80 (m, 3H), 2.93 (m, 1H), 3.17 (q, 2H), 4.34 (m, 2H), 6.85-7.30 (m, 7H), 9.26 (s, 1H), 12.01 (s, 1H).

WO 94/29274

PCT/SE94/00552

-22-

Example 25Preparation of 3-propanoyl-4-(2-methylphenylamino)-8-(3-propylsulfonyl-propoxy)quinoline.

5 3-Propanoyl-4-(2-methylphenylamino)-8-(3-propylthiopropoxy)quinoline
(200 mg, 0.47 mmol) was dissolved in 5 ml methylene chloride. A solution
of sodium bicarbonate (160 mg, 1.90 mmol) in 5 ml water was added. The
mixture was cooled to 2-4°C. A solution of 70 % m-CPBA (230 mg, 0.94
mmol) in 5 ml methylene chloride was added dropwise during 5 min. The
10 temperature was allowed to rise and the mixture was stirred for 30 min at
room temperature. The methylene chloride layer was separated and
washed with water. The organic layer was then dried over Na₂SO₄ and
evaporated. The crude product was purified by column chromatography
(SiO₂; methylene chloride : ethyl acetate 50:50). 110 mg (51%) of the title
15 compound was obtained.

(¹H-NMR, 300 MHz, CDCl₃): 1.06 (t, 3H), 1.28 (t, 3H), 1.88 (m, 2H), 2.35
(s, 3H), 2.50 (m, 2H), 2.97 (t, 2H), 3.16 (q, 2H), 3.32 (t, 2H), 4.35 (t, 2H),
6.85-7.30 (m, 7H), 9.19 (s, 1H), 11.78 (s, 1H).

20

Example 26Preparation of 3-butyryl-4-(4-hydroxy-2-methylphenylamino)-8-(2-methylthioethoxy)quinoline

25 A mixture of 3-butyryl-4-chloro-8-(2-methylthioethoxy)quinoline (2.48 g,
6.63 mmol) and 4-hydroxy-2-methylaniline (1.06 g, 8.62 mmol) in
acetonitrile (20 ml) was refluxed for 8 h. The reaction mixture was
evaporated and the residue was chromatographed (SiO₂; CH₂Cl₂: MeOH
95:5) yielding 1.22 g (45%) of the desired product.

30

(¹H-NMR, 300 MHz, CDCl₃) 1.05 (t, 3H), 1.80 (m, 2H), 2.20 (s, 6 H), 3.00-
3.10 (m, 4H), 4.30 (m, 2H), 6.55 (m, 1H), 6.75-6.85 (m, 2 H), 6.95 (m, 2H),
7.00-7.10 (m, 1 H), 9.15 (s, 1H).

WO 94/29274

PCT/SE94/00552

-23-

Examples 27 and 28Preparation of 3-butyryl-4-(4-hydroxy-2-methylphenylamino)-8-(2-methylsulfinyloethoxy)quinoline (Example 27) and 3-butyryl-4-(4-hydroxy-2-methylphenylamino)-8-(2-methylsulfonyloethoxy)quinoline (Example 28)

5

3-butyryl-4-(4-hydroxy-2-methylphenylamino)-8-(2-methylthioethoxy)-quinoline (1.06 g, 2.59 mmol) was dissolved in methylene chloride (25 ml). 0.3 M NaHCO₃-solution (24 ml) was added. The mixture was cooled to 4°C. A solution of 70% m-CPBA (0.82 g, 3.37 mmol) in methylene chloride (15 ml) was added dropwise. After stirring for 1.5 h at 2-4°C, the organic layer was washed with 0.3 M NaHCO₃-solution. The organic layer was dried over Na₂SO₄ and evaporated. Chromatography on silica gel with EtOAc:CH₂Cl₂ 1:1 as eluent gave 0.4 g (35%) of Example 27 followed by CH₂Cl₂:MeOH 9:1 yielding 0.52 g (47%) of Example 28.

15

Example 27: (¹H-NMR, 300 MHz, CDCl₃) 1.05 (t, 3H), 1.80 (m, 2H), 2.20 (s, 3H), 2.75 (s, 3H), 3.05 (m, 2H), 3.10-3.20 (m, 1H), 3.40-3.50 (m, 1H), 4.55 (m, 2H), 6.55-6.60 (m, 1H), 6.75-6.80 (m, 2H), 6.90-6.95 (m, 1H), 7.00-7.10 (m, 2H), 9.15 (s, 1H).

20

Example 28: (¹H-NMR, 300 MHz, CDCl₃) 1.05 (t, 3H), 1.80 (m, 2H), 2.25 (s, 3H), 3.05 (t, 2H), 3.35 (s, 3H), 3.60 (m, 2H), 4.55 (m, 2H), 6.55-6.60 (m, 1H), 6.75 (m, 1H), 6.80-6.85 (m, 1H), 6.95-7.05 (m, 2H), 7.10 (m, 1H), 9.10 (s, 1H).

25

Example 29Preparation of 3-butyryl-4-(2-chlorophenylamino)-8-(2-methylthioethoxy)quinoline

A mixture of 3-butyryl-4-chloro-8-(2-methylthioethoxy)quinoline (0.8 g, 2.5 mmol) and 2-chloroaniline (0.47 g, 3.7 mmol) in toluene (12 ml) was heated to 90°C and stirred 3.0 h. After cooling to room temperature, methylene chloride and water were added. The mixture was neutralized with a saturated solution of sodium bicarbonate. The organic layer was dried over

30

WO 94/29274

PCT/SE94/00552

-24-

sodium sulfate and evaporated. Trituration with isopropyl ether gave 0.84 g (81 %) of the desired product.

(¹H-NMR, 300 MHz, CDCl₃) : 1.05 (t, 3H), 1.80-1.90 (m, 2H), 2.25 (s, 3H),
5 3.10-3.15 (m, 4H), 4.35-4.40 (m, 2H), 6.80-6.90 (m, 1H), 7.05-7.15 (m, 5H),
7.45-7.50 (m, 1H), 9.30 (s, 1H), 11.55 (s, 1H)

Example 30

Preparation of 3-butyryl-4-(2-chlorophenylamino)-8-(2-methylsulfinylethoxy)quinoline

3-butyryl-4-(2-chlorophenylamino)-8-(2-methylthioethoxy)quinoline (0.34g, 0.82 mmol) was dissolved in methylene chloride (4 ml). Water (2 ml) and sodium hypochlorite (5% in water) (1.37 ml) were added and the mixture
15 was stirred for 2 h. Another portion of sodium hypochlorite (0.5 ml) was added and the stirring was continued for 2 h. The organic layer was dried over sodium sulfate and evaporated. The residue crystallized from a mixture of ethyl acetate and isopropyl ether and 0.25 g (71%) of the title compound was obtained.

20 (¹H-NMR, 300 MHz, CDCl₃) 1.05 (t, 3H), 1.75-1.90 (m, 2H), 2.85 (s, 3H),
3.10 (t, 2H), 3.20-3.30 (m, 1H), 3.45-3.55 (m, 1H), 4.60-4.70 (m, 2H), 6.80-6.90 (m, 1H), 7.00-7.20 (m, 5H), 7.40-7.50 (m, 1H), 9.30 (s, 1H), 11.60 (s, 1H)

Example 31

Preparation of 3-butyryl-4-(2-chlorophenylamino)-8-(2-methylsulfonylethoxy)quinoline

A mixture of 3-butyryl-4-(2-chlorophenylamino)-8-(2-methylthioethoxy)quinoline (0.4 g, 0.96 mmol) in methylene chloride (5 ml) and a saturated solution of sodium bicarbonate (5 ml) was cooled to 4°C. A solution of 70% m-CPBA (0.48 g, 1.97 mmol) in methylene chloride (5 ml) was added dropwise. After stirring for 1 h at 4°C, the organic layer was
30

WO 94/29274

PCT/SE94/00552

-25-

washed with a saturated solution of sodium bicarbonate and thereafter dried over sodium sulfate and evaporated. Trituration with isopropyl ether gave 0.28 g (65%) of the desired product.

5 (¹H-NMR, 300 MHz, CDCl₃) 1.05 (t, 3H), 1.75-1.90 (m, 2H), 3.10 (t, 2H), 3.40 (s, 3H), 3.60-3.70 (m, 2H), 4.60-4.70 (m, 2H), 6.85-6.90 (m, 1H), 7.00-7.20 (m, 5 H), 7.45-7.50 (m, 1H), 9.20 (s, 1H), 11.65 (s, 1H)

Example 32

10 Preparation of 3-butyryl-4-(2-methoxyphenylamino)-8-(2-methylthioethoxy)quinoline

A mixture of 3-butyryl-4-chloro-8-(2-methylthioethoxy)quinoline (0.8 g, 2.5 mmol) and 2-methoxyaniline (0.45 g, 3.7 mmol) in toluene (12 ml) was
15 heated to 90°C and stirred 3.0 h. After cooling to room temperature, methylene chloride and water were added. The mixture was neutralized with a saturated solution of sodium bicarbonate. The organic layer was dried over sodium sulfate and evaporated. Trituration with isopropyl ether gave 0.80 g (77%) of the desired product.

20 (¹H-NMR, 300 MHz, CDCl₃) 1.05 (t, 3H), 1.75-1.90 (m, 2H), 2.30 (s, 3H), 3.05-3.15 (m, 4H), 3.85 (s, 3H), 4.35-4.40 (m, 2H), 6.75-6.85 (m, 1H), 6.90-7.15 (m, 5 H), 7.25-7.30 (m, 1H), 9.25 (s, 1H), 11.55 (s, 1H)

25 Example 33

Preparation of 3-butyryl-4-(2-methoxyphenylamino)-8-(2-methylsulfinylethoxy)quinoline

3-butyryl-4-(2-methoxyphenylamino)-8-(2-methylthioethoxy)quinoline
30 (0.35g, 0.85 mmol) was dissolved in methylene chloride (4 ml). Water (2 ml) and sodium hypochlorite (5% in water) (1.42 ml) were added and the mixture was stirred for 2 h. Another portion of sodium hypochlorite (0.5 ml) was added and the stirring was continued for 2 h. The organic layer

WO 94/29274

PCT/SE94/00552

-26-

was dried over sodium sulfate and evaporated. The residue crystallized from a mixture of ethyl acetate and isopropyl ether and 0.32 g (88 %) of the title compound was obtained.

- 5 (¹H-NMR, 300 MHz, CDCl₃) : 1.05 (t, 3H), 1.75-1.90 (m, 2H), 2.80 (s, 3H), 3.05-3.10 (m, 2H), 3.20-3.30 (m, 1H), 3.45-3.55 (m, 1H), 3.80 (s, 3H), 4.60-4.70 (m, 2H), 6.80-6.85 (m, 1H), 6.90-7.20 (m, 5H), 7.30-7.35 (m, 1H), 9.20 (s, 1H), 11.60 (s, 1H)

10 Example 34

Preparation of 3-butyryl-4-(2-methoxyphenylamino)-8-(2-methylsulfonylethoxy)quinoline

- A mixture of 3-butyryl-4-(2-methoxyphenylamino)-8-(2-methylthioethoxy)quinoline (0.35 g, 0.85 mmol) in methylene chloride (5 ml) and a saturated solution of sodium bicarbonate (5 ml) was cooled to 4°C. A solution of 70% m-CPBA (0.43 g, 1.74 mmol) in methylene chloride (5 ml) was added dropwise. After stirring for 1 h at 4°C, the organic layer was washed with a saturated solution of sodium bicarbonate and thereafter
- 15 dried over sodium sulfate and evaporated. Trituration with isopropyl ether gave 0.28 g (65%) of the desired product.
- 20

- (¹H-NMR, 300 MHz, CDCl₃) : 1.05 (t, 3H), 1.75-1.90 (m, 2H), 3.05-3.10 (m, 2H), 3.40 (s, 3H), 3.60-3.70 (m, 2H), 3.85 (s, 3H), 4.60-4.65 (m, 2H), 6.80-6.85
- 25 (m, 1H), 6.90-7.20 (m, 5H), 7.30-7.35 (m, 1H), 9.15 (s, 1H), 11.60 (s, 1H)

Example 35

Preparation of 3-butyryl-4-(2,4-dimethylphenylamino)-8-(2-methylthioethoxy)quinoline

- 30 A mixture of 3-butyryl-4-chloro-8-(2-methylthioethoxy)quinoline (0.8 g, 2.5 mmol) and 2,4-dimethylaniline (0.45 g, 3.7 mmol) in toluene (12 ml) was heated to 90°C and stirred 3.0 h. After cooling to room temperature,

WO 94/29274

PCT/SE94/00552

-27-

methylene chloride and water were added. The mixture was neutralized with a saturated solution of sodium bicarbonate. The organic layer was dried over sodium sulfate and evaporated. Trituration with isopropyl ether gave 0.77 g (75 %) of the desired product.

5

$^1\text{H-NMR}$, 300 MHz, CDCl_3) : 1.05 (t, 3H), 1.75-1.90 (m, 2H), 2.25 (s, 3H), 2.30 (s, 3H), 2.35 (s, 3H), 3.05-3.15 (m, 4H), 4.30-4.40 (m, 2H), 6.80-6.85 (m, 1H), 6.90-7.10 (m, 5H), 9.20 (s, 1H), 11.85 (s, 1H)

10 Example 36

Preparation of 3-butyryl-4-(2,4-dimethylphenylamino)-8-(2-methylsulfinylethoxy)quinoline

3-butyryl-4-(2,4-dimethylphenylamino)-8-(2-methylthioethoxy)quinoline (0.34g, 0.83 mmol) was dissolved in methylene chloride (4 ml). Water (2 ml) and sodium hypochlorite (5% in water) (1.39 ml) were added and the mixture was stirred for 2 h. Another portion of sodium hypochlorite (0.5 ml) was added and the stirring was continued for 2 h. The organic layer was dried over sodium sulfate and evaporated. Trituration with isopropyl ether gave 0.32 g (91 %) of the title compound.

20

$^1\text{H-NMR}$, 300 MHz, CDCl_3) 1.05 (t, 3H), 1.75-1.90 (m, 2H), 2.30 (s, 3H), 2.35 (s, 3H), 2.80 (s, 3H), 3.05-3.15 (m, 2H), 3.20-3.30 (m, 1H), 3.40-3.55 (m, 1H), 4.60-4.65 (m, 2H), 6.80-6.85 (m, 1H), 6.90-7.15 (m, 5H), 9.15 (s, 1H), 11.85 (s, 1H)

25

Example 37

Preparation of 3-butyryl-4-(2,4-dimethylphenylamino)-8-(2-methylsulfonylethoxy)quinoline

30 A mixture of 3-butyryl-4-(2,4-dimethylphenylamino)-8-(2-methylthioethoxy)quinoline (0.33 g, 0.81 mmol) in methylene chloride (5 ml) and a saturated solution of sodium bicarbonate (5 ml) was cooled to 4°C. A solution of 70% m-CPBA (0.40 g, 1.66 mmol) in methylene chloride

WO 94/29274

PCT/SE94/00552

-28-

(5 ml) was added dropwise. After stirring for 1 h at 4°C, the organic layer was washed with a saturated solution of sodium bicarbonate and thereafter dried over sodium sulfate and evaporated. After trituration with isopropyl ether a crystalline product was obtained. Chromatography (SiO₂; CH₂Cl₂: MeOH 90:10) gave 0.17 g (48%) of the desired compound.

(¹H-NMR, 300 MHz, CDCl₃) 1.05 (t, 3H), 1.75-1.90 (m, 2H), 2.30 (s, 3H), 2.35 (s, 3H), 3.05-3.15 (m, 2H), 3.40 (s, 3H), 3.60-3.65 (m, 2H), 4.60-4.65 (m, 2H), 6.80-6.85 (m, 1H), 6.90-7.15 (m, 5H), 9.10 (s, 1H), 11.90 (s, 1H)

10

Example 38

Preparation of 3-butyryl-4-(2,6-dimethylphenylamino)-8-(2-methylthioethoxy)quinoline

15 A mixture of 3-butyryl-4-chloro-8-(2-methylthioethoxy)quinoline (0.8 g, 2.5 mmol) and 2,6-dimethylaniline (0.45 g, 3.7 mmol) in toluene (12 ml) was heated to 90°C and stirred 3.0 h. After cooling to room temperature, methylene chloride and water were added. The mixture was neutralized with a saturated solution of sodium bicarbonate. The organic layer was
20 dried over sodium sulfate and evaporated. Chromatography (SiO₂; EtOAc) gave 0.7 g (68 %) of the title compound.

(¹H-NMR, 300 MHz, CDCl₃) 1.05 (t, 3H), 1.75-1.90 (m, 2H), 2.10 (s, 6H), 2.25 (s, 3H), 3.05-3.15 (m, 4H), 4.30-4.35 (m, 2H), 6.85-7.20 (m, 6H), 9.20 (s, 1H), 12.25 (s, 1H)

25

Example 39

Preparation of 3-butyryl-4-(2,6-dimethylphenylamino)-8-(2-methylsulfinylethoxy)quinoline

30

3-butyryl-4-(2,6-dimethylphenylamino)-8-(2-methylthioethoxy)quinoline (0.33 g, 0.81 mmol) was dissolved in methylene chloride (4 ml). Water (2 ml) and sodium hypochlorite (5% in water) (1.7 ml) were added and the

WO 94/29274

PCT/SE94/00552

-29-

mixture was stirred for 3 h. The organic layer was dried over sodium sulfate and evaporated. Trituration with isopropyl ether gave 0.20 g (58 %) of the title compound.

5 (¹H-NMR, 300 MHz, CDCl₃) 1.05 (t, 3H), 1.80-1.90 (m, 2H), 2.10 (s, 3H), 2.15 (s, 3H), 2.80 (s, 3H), 3.10-3.15 (m, 2H), 3.20-3.30 (m, 1H), 3.40-3.55 (m, 1H), 4.55-4.65 (m, 2H), 6.85-6.95 (m, 2H), 7.05-7.25 (m, 4H), 9.20 (s, 1H), 12.25 (s, 1H)

10 Example 40

Preparation of 3-butyryl-4-(2,6-dimethylphenylamino)-8-(2-methylsulfonylethoxy)quinoline

A mixture of 3-butyryl-4-(2,6-dimethylphenylamino)-8-(2-methylthioethoxy)quinoline (0.36 g, 0.88 mmol) in methylene chloride (5 ml) and a saturated solution of sodium bicarbonate (5 ml) was cooled to 4°C. A solution of 70% m-CPBA (0.42 g, 1.76 mmol) in methylene chloride (5 ml) was added dropwise. After stirring for 1 h at 4°C, the organic layer was washed with a saturated sodium bicarbonate solution and thereafter
15 dried over sodium sulfate and evaporated. After trituration with isopropyl ether a crystalline product was obtained. Chromatography (SiO₂; CH₂Cl₂: MeOH 90:10) and finally crystallization from ethyl acetate gave 0.070 g (18%) of the desired product.

25 (¹H-NMR, 300 MHz, CDCl₃) 1.05 (t, 3H), 1.80-1.95 (m, 2H), 2.10 (s, 6H), 3.05-3.15 (m, 2H), 3.40 (s, 3H), 3.60-3.65 (m, 2H), 4.55-4.60 (m, 2H), 6.85-6.95 (m, 2H), 7.00-7.25 (m, 4H), 9.10 (s, 1H), 12.30 (s, 1H)

WO 94/29274

PCT/SE94/00552

-30-

Example 41Preparation of 3-butyryl-4-(2-methyl,6-chlorophenylamino)-8-(2-methylthioethoxy)quinoline

5 A mixture of 3-butyryl-4-chloro-8-(2-methylthioethoxy)quinoline (0.8 g, 2.5 mmol) and 2-methyl, 6-chloroaniline (0.52 g, 3.7 mmol) in toluene (12 ml) was heated to 90°C and stirred 3.0 h. After cooling to room temperature, methylene chloride and water were added. The mixture was neutralized with a saturated solution of sodium bicarbonate. The organic layer was
10 dried over sodium sulfate and evaporated. Chromatography (SiO₂; EtOAc) gave 0.77 g (72 %) of the title compound.

(¹H-NMR, 300 MHz, CDCl₃) 1.05 (t, 3H), 1.80-1.90 (m, 2H), 2.15 (s, 3H), 2.25 (s, 3H), 3.05-3.15 (m, 4H), 4.30-4.40 (m, 2H), 6.85-6.90 (1H), 6.90-7.05
15 (m, 2H), 7.15-7.35 (m, 3H), 9.20 (s, 1H), 12.15 (s, 1H)

Example 42Preparation of 3-butyryl-4-(2-methyl,6-chlorophenylamino)-8-(2-methylsulfinylethoxy)quinoline

20 3-butyryl-4-(2-methyl,6-chlorophenylamino)-8-(2-methylthioethoxy)quinoline (0.35 g, 0.82 mmol) was dissolved in methylene chloride (4 ml). Water (2 ml) and sodium hypochlorite (5% in water) (1.7 ml) were added and the mixture was stirred for 3 h. The organic layer was dried over
25 sodium sulfate and evaporated. The residue crystallized from ethyl acetate and 0.10 g (27%) of the title compound was obtained.

(¹H-NMR, 300 MHz, CDCl₃) 1.05 (t, 3H), 1.80-1.90 (m, 2H), 2.15 (m, 3H), 2.85 (m, 3H), 3.10-3.15 (m, 2H), 3.20-3.30 (m, 1H), 3.40-3.55 (m, 1H), 4.55-
30 4.70 (m, 2H), 6.85-7.00 (m, 2H), 7.05-7.10 (m, 1H), 7.15-7.35 (m, 3H), 9.20 (s, 1H), 12.20 (s, 1H)

WO 94/29274

PCT/SE94/00552

-31-

Example 43Preparation of 3-butyryl-4-(2-methyl,6-chlorophenylamino)-8-(2-methylsulfonylethoxy)quinoline

5 A mixture of 3-butyryl-4-(2-methyl,6-chlorophenylamino)-8-(2-methylthioethoxy)quinoline (0.34 g, 0.79 mmol) in methylene chloride (5 ml) and a saturated solution of sodium bicarbonate (5 ml) was cooled to 4°C. A solution of 70% m-CPBA (0.38 g, 1.58 mmol) in methylene chloride (5 ml) was added dropwise. After stirring for 1 h at 4°C, the organic layer
10 was washed with a saturated sodium bicarbonate solution and thereafter dried over sodium sulfate and evaporated. Crystallization from ethyl acetate gave 0.11 g (30%) of the desired product.

(¹H-NMR, 300 MHz, CDCl₃) 1.05 (t, 3H), 1.80-1.95 (m, 2H), 2.15 (s, 3H),
15 3.05-3.15 (m, 2H), 3.40 (s, 3H), 3.60-3.70 (m, 2H), 4.55-4.65 (m, 2H), 6.90-7.05 (m, 3H), 7.15-7.25 (m, 2H), 7.30-7.35 (m, 1H), 9.15 (s, 1H), 12.20 (s, 1H)

Example 44Preparation of 3-propanoyl-4-(2-methylphenylamino)-8-(3-methylthiopropoxy)quinoline.

20 3-Propanoyl-4-chloro-8-(3-methylthiopropoxy)quinoline (1.2 g, 3.71 mmol) and o-toluidine (0.795 g, 7.42 mmol) was refluxed in acetonitrile (18 ml) for 100 min. The solvent was evaporated and the residue was purified by
25 column chromatography (methylene chloride : methanol 100:3). 1.45 g (99%) of the title compound was obtained.

(¹H-NMR, 300 MHz, CDCl₃) 1.3 (t, 3H) , 2.15 (s, 3H), 2.2-2.3 (m, 2H), 2.33 (s, 3H), 2.75 (t, 2H), 3.15 (q, 2H), 4.28 (t, 2H), 6.83-6.92 (m, 1H), 6.95-7.18
30 (m, 5H), 7.25-7.33 (m, 1H), 9.25 (s, 1H), 11.75 (s, 1H).

WO 94/29274

PCT/SE94/00552

-32-

Examples 45 and 46Preparation of 3-propanoyl-4-(2-methylphenylamino)-8-(3-methylsulfinylpropoxy)quinoline (Example 45) and 3-propanoyl-4-(2-methylphenylamino)-8-(3-methylsulfonylpropoxy)quinoline (Example 46).

5

3-Propanoyl-4-(2-methylphenylamino)-8-(3-methylthiopropoxy) quinoline (1.03 g, 2.611 mmol) was dissolved in methylene chloride (30 ml). 0.3 M NaHCO₃ solution (26 ml, 7.83 mmol) was added. The mixture was cooled to 4°C. A solution of 70.5% m-CPBA (0.895 g, 3.66 mmol) in methylene chloride (27 ml) was added dropwise over a 45 min period. After stirring for 30 min at 4°C, the organic layer was separated and washed with 0.3 M NaHCO₃ solution. The organic layer was dried over Na₂SO₄ and evaporated. Column chromatography (SiO₂; CH₂Cl₂:MeOH 100:3 and 100:6) gave 0.48 g (45%) of the compound according to Example 45 and 0.50 g (45%) of the compound according to Example 46.

Example 45: (¹H-NMR, 300 MHz, CDCl₃) 1.3 (t, 3H), 2.38 (s, 3H), 2.44-2.55 (m, 2H), 2.65 (s, 3H), 2.9-3.02 (m, 1H), 3.08-3.23 (m, 3H), 4.28-4.4 (m, 2H), 6.85-6.92 (m, 1H), 6.95-7.18 (m, 5H), 7.25-7.34 (m, 1H), 9.25 (s, 1H), 11.8 (s, 1H)

Example 46: (¹H-NMR, 300 MHz, CDCl₃) 1.28 (t, 3H), 2.35 (s, 3H), 2.45-2.58 (m, 2H), 2.95 (s, 3H), 3.18 (q, 2H), 3.4 (t, 2H), 4.35 (t, 2H), 6.85-6.92 (m, 1H), 6.95-7.15 (m, 5H), 7.22-7.33 (m, 1H), 9.23 (s, 1H), 11.83 (s, 1H)

25

Example 47Preparation of 3-butyryl-4-(2-chlorophenylamino)-8-(3-methylthiopropoxy)quinoline.

30 A mixture of 3-butyryl-4-chloro-8-(3-methylthiopropoxy)quinoline (0.95 g, 2.8 mmol) and 2-chloroaniline (1.51 g, 11.8 mmol) in toluene was heated to 55°C and stirred overnight. The solvent was evaporated and the residue was partitioned between methylene chloride and a saturated sodium

WO 94/29274

PCT/SE94/00552

-33-

bicarbonate solution. The organic layer was dried over Na_2SO_4 and evaporated. The residue was chromatographed (SiO_2 ; CH_2Cl_2 :MeOH 95:5) yielding 0.95 g (74%) of the desired product.

- 5 (^1H -NMR, 500 MHz, CDCl_3) 0.95-1.05 (t, 3H), 1.75-1.85 (m, 2H), 2.10 (s, 3H), 2.20-2.30 (m, 2H), 2.70-2.80 (m, 2H), 3.0-3.10 (m, 2H), 4.20-4.30 (m, 2H), 6.80 (d, 1H), 6.95-7.10 (m, 5H), 7.40 (d, 1H), 9.20 (s, 1H), 11.6 (s, 1H).

Example 48

- 10 Preparation of 3-butyryl-4-(2-chlorophenylamino)-8(3-methylsulfinylpropoxy)quinoline.

3-butyryl-4-(2-chlorophenylamino)-8-(3-methylthiopropoxy)quinoline (0.31 g, 0.72 mmol) was dissolved in methylene chloride (10ml), 5 ml water was added and then a solution of 1.5 ml (1.09 mmol) of 5% NaOCl in 10 ml methylene chloride was added. The mixture was stirred for 4 h at room temperature. The organic phase was separated and evaporated. Chromatography with methylene chloride : methanol 95:5 as the eluent gave 0.104g (32%) of the title compound.

- 20 (^1H -NMR, 500 MHz, CDCl_3) 1.05 (t, 3H), 1.80-1.90 (m, 3H), 2.45-2.55 (m, 2H), 2.65 (s, 3H), 2.90-3.0 (m, 1H), 3.05-3.15 (m, 2H), 3.15-3.20 (m, 1H), 4.30-4.40 (m, 2H), 6.85 (d, 1H), 7.0-7.15 (m, 5H), 7.45 (d, 1H), 9.25 (s, 1H), 11.60 (s, 1H)

25

Example 49

Preparation of 3-butyryl-4-(2-chlorophenylamino)-8(3-methylsulfonylpropoxy)quinoline.

- 30 A mixture of 3-butyryl-4(2-chlorophenylamino)-8-(3-methylthiopropoxy)quinoline (0.31 g, 0.72 mmol) in 5 ml methylene chloride and NaHCO_3 (0.27 g, 3.2 mmol) in 5 ml H_2O was cooled to 4°C . A solution of 70% m-CPBA (0.4 g, 1.63 mmol) in 10 ml methylene chloride

WO 94/29274

PCT/SE94/00552

-34-

was added dropwise. After stirring for 1 h at 4°C, the organic layer was washed with a saturated sodium bicarbonate solution. The organic layer was dried over Na₂SO₄ and evaporated. Chromatography (SiO₂; CH₂Cl₂:MeOH 95:5) gave 69 mg (21%) of the desired product.

5

(¹H-NMR, 500 MHz, CDCl₃) 1.05 (t, 3H), 1.80-1.90 (m, 2H), 2.50-2.55 (m, 2H), 3.00 (s, 3H), 3.05-3.10 (m, 2H), 3.40-3.45 (m, 2H), 4.35-4.40 (m, 2H), 6.80-6.85 (m, 1H), 7.00-7.20 (m, 5H), 7.45-7.50 (m, 1H), 9.25 (s, 1H), 11.60 (s, 1H)

10

Example 50

Preparation of 3-butyryl-4-(2-methylphenylamino)-8-(3-methylthiopropoxy)quinoline.

15 A mixture of 3-butyryl-4-chloro-8-(3-methylthiopropoxy)quinoline (0.95 g, 2.97 mmol) and 2-methylaniline (1.27 g, 11.8 mmol) in toluene was heated to 55°C and stirred overnight. The solvent was evaporated and the residue was partitioned between methylene chloride and a saturated sodium bicarbonate solution. The organic layer was dried over Na₂SO₄ and
20 evaporated. The residue was chromatographed (SiO₂ ; CH₂Cl₂:MeOH 95:5) yielding 0.98 g (80.9%) of the desired product.

25 (¹H-NMR, 500 MHz, CDCl₃) 1.05 (t, 3H), 1.75-1.80 (m, 2H), 2.10 (s, 3H), 2.25-2.30 (m, 2H), 2.35 (s, 3H), 2.75-2.80 (m, 2H), 3.05-3.10 (m, 2H), 4.25-4.30 (m, 2H), 7.85-7.90 (d, 1H), 6.95-7.15 (m, 5H), 7.25 (d, 1H), 9.20 (s, 1H), 11.75 (s, 1H)

WO 94/29274

PCT/SE94/00552

-35-

Example 51Preparation of 3-butyryl-4-(2-methylphenylamino)-8-(3-methylsulfinylpropoxy)quinoline.

- 5 3-butyryl-4-(2-methylphenylamino)-8-(3-methylthiopropoxy)quinoline (0.33 g, 0.8 mmol) was dissolved in methylene chloride (15 ml), 5 ml water was added and then a solution of 1.5 ml (1.09 mmol) of 5% NaOCl in 10 ml methylene chloride was added. The mixture was stirred for 4 h at room temperature. The organic phase was separated and evaporated.
- 10 Chromatography with methylene chloride : methanol 95:5 as the eluent gave 0.18g (53%) of the title compound.

- (¹H-NMR, 500 MHz, CDCl₃) 1.05 (t, 3H), 1.75-1.85 (m, 2H), 2.30 (s, 3H), 2.40-2.50 (m, 2H), 2.60 (s, 3H), 2.90-3.15 (m, 4H), 4.25-4.40 (m, 2H), 6.85-6.90 (m, 1H), 6.95-7.15 (m, 5H), 7.25-7.30 (m, 1H), 9.20 (s, 1H), 11.90 (s, 1H)
- 15

Example 52Preparation of 3-butyryl-4-(2-methylphenylamino)-8-(3-methylsulfonylpropoxy)quinoline.

- 20 A mixture of 3-butyryl-4(2-methylphenylamino)-8-(3-methylthiopropoxy)quinoline (0.33 g, 0.81 mmol) in 5 ml methylene chloride and NaHCO₃ (0.27 g, 3.2 mmol) in 5 ml H₂O was cooled to 4°C. A solution of 70% m-CPBA (0.4 g, 1.63 mmol) in 10 ml methylene chloride
- 25 was added dropwise. After stirring for 1 h at 4°C, the organic layer was washed with a saturated sodium bicarbonate solution. The organic layer was dried over Na₂SO₄ and evaporated. Chromatography (SiO₂; CH₂Cl₂:MeOH 95:5) gave 99 mg (28%) of the desired product.

- 30 (¹H-NMR, 500 MHz, CDCl₃) : 1.05 (t, 3H), 1.80-1.90 (m, 2H), 2.35 (s, 3H), 2.50-2.55 (m, 2H), 3.00 (s, 3H), 3.10-3.15 (m, 2H), 3.35-3.45 (m, 2H), 4.35-4.40 (m, 2H), 6.85-6.90 (m, 1H), 6.95-7.15 (m, 5H), 7.25-7.30 (m, 1H), 9.20 (s, 1H), 11.85 (s, 1H)

WO 94/29274

PCT/SE94/00552

-36-

Example 53Preparation of 3-butyryl-4-(2-methoxyphenylamino)-8-(3-methylthiopropoxy)quinoline.

5 A mixture of 3-butyryl-4-chloro-8-(3-methylthiopropoxy)quinoline (0.95 g, 2.97 mmol) and 2-methoxyaniline (1.46 g, 11.8 mmol) in toluene was heated to 55°C and stirred overnight. The solvent was evaporated and the residue was partitioned between methylene chloride and a saturated sodium bicarbonate solution. The organic layer was dried over Na₂SO₄ and evaporated. The residue was chromatographed (SiO₂ ; CH₂Cl₂:MeOH 95:5) yielding 0.90g (71%) of the desired product.

10 (¹H-NMR, 300 MHz, CDCl₃) : 1.05 (t, 3H), 1.75-1.90 (m, 2H), 2.10 (s, 3H), 2.25-2.35 (m, 2H), 2.75-2.80 (m, 2H), 3.05-3.10 (m, 2H), 3.80 (s, 3H), 4.25-4.35 (m, 2H), 6.75-6.85 (m, 1H), 6.90-7.20 (m, 5H), 7.25-7.30 (m, 1H), 9.20 (s, 1H), 11.65 (s, 1H)

Example 54

20 Preparation of 3-butyryl-4-(2-methoxyphenylamino)-8-(3-methylsulfinylpropoxy)quinoline.

3-butyryl-4-(2-methoxyphenylamino)-8-(3-methylthiopropoxy)quinoline (0.30 g, 0.70 mmol) was dissolved in methylene chloride (10 ml), 5 ml water was added and then a solution of 1.5 ml (1.09 mmol) of 5% NaOCl in 10 ml methylene chloride was added. The mixture was stirred for 4 h at room temperature. The organic phase was separated and evaporated. Chromatography with methylene chloride : methanol 95:5 as the eluent gave 14 mg (4.6%) of the title compound.

30 (¹H-NMR, 500 MHz, CDCl₃) 1.0 (t, 3H), 1.75-1.85 (m, 2H), 2.45-2.55 (m, 2H), 2.63 (s, 3H), 2.95 (m, 1H), 3.05-3.10 (m, 2H), 3.15 (m, 1H), 3.8 (s, 3H), 4.30 (m, 1H), 4.40 (m, 1H), 6.80-7.20 (m, 6H), 7.30 (m, 1H), 9.20 (s, 1H), 11.6 (s, 1H)

WO 94/29274

PCT/SE94/00552

-37-

Example 55Preparation of 3-butyryl-4-(2-methoxyphenylamino)-8-(3-methylsulfonylpropoxy)quinoline.

5 A mixture of 3-butyryl-4-(2-methoxyphenylamino)-8-(3-methylthiopropoxy)quinoline (0.30 g, 0.71 mmol) in 5 ml methylene chloride and NaHCO_3 (0.27 g, 3.2 mmol) in 5 ml H_2O was cooled to 4°C . A solution of 70% m-CPBA (0.4 g, 1.63 mmol) in 10 ml methylene chloride was added dropwise. After stirring for 1 h at 4°C , the organic layer was
10 washed with a saturated sodium bicarbonate solution. The organic layer was dried over Na_2SO_4 and evaporated. Chromatography (SiO_2 ; CH_2Cl_2 :MeOH 95:5) gave 41 mg (13%) of the desired product.

(^1H -NMR, 500 MHz, CDCl_3) 1.05 (t, 3H), 1.80-1.90 (m, 2H), 2.50-2.55 (m, 2H), 3.00 (s, 3H), 3.05-3.10 (m, 2H), 3.40-3.45 (m, 2H), 3.80 (s, 3H),
15 4.35-4.40 (m, 2H), 6.80-7.15 (m, 6H), 7.30-7.35 (m, 1H), 9.20 (s, 1H), 11.60 (s, 1H)

Example 56

20 Resolution of 3-butyryl-4-(2-methylphenylamino)-8-(2-methylsulfinylethoxy)quinoline

A mixture of 3-butyryl-4-(2-methylphenylamino)-8-(2-methylsulfinylethoxy)quinoline (9.3 g, 0.023 mmol) and D-(-)-tartaric acid
25 (3.45 g, 0.023 mmol) in methanol (180 ml) was heated to reflux. The solution was allowed to cool to room temperature and stirred for 60 h. The precipitate was filtered off and washed with a total amount of 20 ml methanol giving 6.1 g of the tartaric salt (the filtrate was used in Example 57). Recrystallization from methanol was repeated 3 times yielding 3.05 g,
30 1.30 g and finally 1.05 g of the tartaric salt of Example 57. The salt was neutralized with a saturated sodium bicarbonate in methylene chloride and water. The organic layer was dried over sodium sulfate and the solvent

WO 94/29274

PCT/SE94/00552

—38—

was evaporated. Trituration with isopropyl ether gave 0.7 g of the pure enantiomer.

Example 57

5 Resolution of 3-butyryl-4-(2-methylphenylamino)-8-(2-methylsulfinylethoxy)quinoline

The filtrate from the first crystallization in Example 56 was evaporated. The salt was neutralized with a saturated sodium bicarbonate solution in methylene chloride and water. The organic layer was dried over sodium sulfate and the solvent was evaporated. The solid residue (4.6 g, 0.011 mole) and L-(+)-tartaric acid (1.68 g, 0.011 mole) were dissolved in warm methanol (110 ml). The solution was allowed to cool to room temperature and stirred for 72 h. The precipitate was filtered off and washed with a total amount of 11 ml methanol giving 1.5 g of the tartaric salt. Recrystallization from methanol gave 1.05 g of the tartaric salt of Example 57. The salt was neutralized with a saturated sodium bicarbonate solution in methylene chloride and water. The organic layer was dried over sodium sulfate and the solvent was evaporated. Trituration with isopropyl ether gave 0.7 g of the pure enantiomer.

The enantiomers were separated on a 250 X 4.6 mm i.d. Chiralpak AD column (Dacel, Japan) using the following parameters:
n-hexane: 2-propanol: acetonitrile: diethyl amine (82: 18: 2: 0.1);
25 temperature: 35°C; flow rate: 0.8 ml/min.

Enantiomer according to Example 56: retention time 14.5 min
Enantiomer according to Example 57: retention time 18.4 min

WO 94/29274

PCT/SE94/00552

-39-

TABLE 1

Summary of the exemplified compounds of the invention.

Example	R ¹	R ²	R ³	R ⁴	m	n
1	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	H	2	0
2	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	H	2	1
3	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	H	2	2
4	CH ₂ CH ₂ CH ₃	CH(CH ₃) ₂	CH ₃	H	2	0
5	CH ₂ CH ₂ CH ₃	CH(CH ₃) ₂	CH ₃	H	2	1
6	CH ₂ CH ₂ CH ₃	CH(CH ₃) ₂	CH ₃	H	2	2
7	CH ₂ CH ₃	CH ₃	CH ₃	H	2	0
8	CH ₂ CH ₃	CH ₃	CH ₃	H	2	1
9	CH ₂ CH ₃	CH ₃	CH ₃	H	2	2
10	CH ₂ CH ₃	CH ₂ CH ₃	CH ₃	H	2	0
11	CH ₂ CH ₃	CH ₂ CH ₃	CH ₃	H	2	1
12	CH ₂ CH ₃	CH ₂ CH ₃	CH ₃	H	2	2
13	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	4-F	2	0
14	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	4-F	2	1
15	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	4-F	2	2
16	CH ₂ CH ₃	CH(CH ₃) ₂	CH ₃	H	2	0
17	CH ₂ CH ₂ CH ₃	CH ₂ CH ₃	CH ₃	H	2	0
18	CH ₂ CH ₂ CH ₃	CH ₂ CH ₃	CH ₃	H	2	1
19	CH ₂ CH ₂ CH ₃	CH ₂ CH ₃	CH ₃	H	2	2
20	CH ₂ CH ₃	CH ₃	CH ₂ CH ₂ CH ₃	H	2	0
21	CH ₂ CH ₃	CH ₃	CH ₂ CH ₂ CH ₃	H	2	1
22	CH ₂ CH ₃	CH ₃	CH ₂ CH ₂ CH ₃	H	2	2
23	CH ₂ CH ₃	CH ₃	CH ₂ CH ₂ CH ₃	H	3	0

WO 94/29274

PCT/SE94/00552

-40-

Example	R ¹	R ²	R ³	R ⁴	m	n
24	CH ₂ CH ₃	CH ₃	CH ₂ CH ₂ CH ₃	H	3	1
25	CH ₂ CH ₃	CH ₃	CH ₂ CH ₂ CH ₃	H	3	2
26	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	4-OH	2	0
27	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	4-OH	2	1
28	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	4-OH	2	2
29	CH ₂ CH ₂ CH ₃	Cl	CH ₃	H	2	0
30	CH ₂ CH ₂ CH ₃	Cl	CH ₃	H	2	1
31	CH ₂ CH ₂ CH ₃	Cl	CH ₃	H	2	2
32	CH ₂ CH ₂ CH ₃	OCH ₃	CH ₃	H	2	0
33	CH ₂ CH ₂ CH ₃	OCH ₃	CH ₃	H	2	1
34	CH ₂ CH ₂ CH ₃	OCH ₃	CH ₃	H	2	2
35	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	4-CH ₃	2	0
36	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	4-CH ₃	2	1
37	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	4-CH ₃	2	2
38	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	6-CH ₃	2	0
39	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	6-CH ₃	2	1
40	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	6-CH ₃	2	2
41	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	6-Cl	2	0
42	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	6-Cl	2	1
43	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	6-Cl	2	2
44	CH ₂ CH ₃	CH ₃	CH ₃	H	3	0
45	CH ₂ CH ₃	CH ₃	CH ₃	H	3	1
46	CH ₂ CH ₃	CH ₃	CH ₃	H	3	2
47	CH ₂ CH ₂ CH ₃	Cl	CH ₃	H	3	0
48	CH ₂ CH ₂ CH ₃	Cl	CH ₃	H	3	1

WO 94/29274

PCT/SE94/00552

-41-

Example	R ¹	R ²	R ³	R ⁴	m	n
49	CH ₂ CH ₂ CH ₃	Cl	CH ₃	H	3	2
50	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	H	3	0
51	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	H	3	1
52	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	H	3	2
53	CH ₂ CH ₂ CH ₃	OCH ₃	CH ₃	H	3	0
54	CH ₂ CH ₂ CH ₃	OCH ₃	CH ₃	H	3	1
55	CH ₂ CH ₂ CH ₃	OCH ₃	CH ₃	H	3	2

WO 94/29274

PCT/SE94/00552

—42—

2. PREPARATION OF INTERMEDIATES

The following examples illustrate intermediates useful in the preparation of compounds according to the invention:

5

Example IPreparation of 2-(2-methylthioethoxy)nitrobenzene

2-methylthioethylchloride (18.0 g, 0.16 mole), o-nitrophenol (20.8 g, 0.15 mole) and potassium carbonate (24.7 g, 0.18 mole) was refluxed in acetonitrile for 24 h. The reaction mixture was filtered and the solvent was evaporated. The residue was dissolved in methylene chloride and washed with water once and thereafter twice with a saturated sodium carbonate solution. The organic layer was dried over sodium sulfate and the solvent was evaporated giving 20.2 g (63%) of the title compound as an oily residue.

15

(¹H-NMR, 300 MHz, CDCl₃), 2.21 (s,3H), 2.90 (t,2H), 4.27 (t,2H), 7.04 (m,2H), 7.50 (m,1H), 7.79 (m,1H).

20

Example IIPreparation of 2-(2-methylthioethoxy)aniline.

Tin chloride dihydrate (57.9 g, 0.26 mole) in ethyl alcohol (90 ml) was added to a mixture of 2-(2-methylthioethoxy)nitrobenzene (18.1g, 0.085 mole), conc. HCl (72.4 ml) and ethyl alcohol (36 ml). The reaction mixture was stirred for 24 h at room temperature. Sodium hydroxide (6 M, 270 ml) was added to the reaction mixture. Extraction with methylene chloride (3 x 400 ml) gave after drying the organic layer over sodium sulfate and evaporation of the solvent 14.8 g (95%) of the title compound.

25

30

(¹H-NMR, 300 MHz, CDCl₃), 2.28 (s,3H), 2.90 (t,2H), 4.18 (t,2H), 4.83 (b,2H), 6.66-6.83 (m,4H).

WO 94/29274

PCT/SE94/00552

-43-

Example IIIPreparation of ethyl 2-butyryl-3-(2-(2-methylthioethoxy)phenylamino)acrylate

5 A mixture of 2-(2-methylthioethoxy)aniline (1.6 g, 8.7 mmol), ethyl butyryl acetate (1.38 g, 8.7 mmol) and triethyl orthoformate (1.30 g, 8.8 mmol) was heated to 120°C for 1 h and ethyl alcohol was distilled off. The reaction mixture was cooled to room temperature. Trituration with methyl alcohol gave 1.08 g (35%) of the desired compound as a solid.

10

$^1\text{H-NMR}$, 300 MHz, CDCl_3) 0.96 (t,3H), 1.33 (t,3H), 1.68 (m,3H), 2.22 (s,3H), 2.92 (t,2H), 3.00 (t,2H), 4.25 (m,4H), 6.9-7.3 (m,4H), 8.5 (d,1H), 12.81 (d,1H).

15

Example IVPreparation of 3-butyryl-8-(2-(2-methylthioethoxy)-4(1H)-quinolone

Ethyl 2-butyryl-4-(2-(2-methylthioethoxy)phenylamino) acrylate (1.07, 3.04 mmol) was added to refluxing diphenyl ether. The mixture was refluxed
20 for 50 min. The reaction mixture was cooled to room temperature. Petroleum ether was added (70 ml) and after stirring the mixture for further 90 min, the precipitate was filtered off giving 0.8 g (85%) of the title compound.

25

$^1\text{H-NMR}$, 500 MHz, CDCl_3) 1.02 (t,3H), 1.75 (m,2H), 2.22 (s,3H), 3.00 (t,2H), 3.25 (t,2H), 4.36 (t,2H), 7.15 (d,1H), 7.35 (m,1H), 8.06 (d,1H), 8.58 (s,1H), 9.40 (b,1H).

WO 94/29274

PCT/SE94/00552

—44—

Example VPreparation of 3-butyryl-4-chloro-8-(2-methylthioethoxy)-quinolone

3-butyryl-8-(2-methylthioethoxy)-4(1H)-quinolone (0.8 g, 2.8 mmol) and
5 phosphorus oxychloride (10 ml) was stirred at room temperature for 1 h.
The phosphorus oxychloride was evaporated. The residue was partitioned
between water and methylene chloride. pH was adjusted to 8 with sodium
bicarbonate. The organic layer was dried over sodium sulfate and the
solvent was evaporated giving 0.57 g (68%) of the desired compound.

10

(¹H-NMR, 300 MHz, CDCl₃), 0.97 (t,3H), 1.86 (m,2H), 2.22 (s,3H), 3.00
(t,2H), 3.05 (t,2H), 4.38 (t,2H), 7.16 (d,1H), 7.57 (m,1H), 7.87 (d,1H), 8.84
(s,1H).

15

Example VIPreparation of 3-propanoyl-4-chloro-8-(2-methylthioethoxy)-quinoline.

The title compound was prepared according to the method in Example V.
Yield: 0.6 g (75%).

20

(¹H-NMR, 300 MHz, CDCl₃) 1.26 (t,3H), 2.26 (s,3H), 3.01-3.07 (m,4H), 4.39
(t,2H), 7.15 (d,1H), 7.55 (m,1H), 7.85 (d,1H), 8.84 (s,1H).

Example VII

25

Preparation of 3-propanoyl-4-chloro-8-(2-propylthioethoxy)-quinoline.

3-Propanoyl-4-chloro-8-(2-propylthioethoxy) quinoline was synthesized
according to Example V. Yield 2.5 g (88%).

30

(¹H-NMR, 300 MHz, CDCl₃) 0.97 (t, 3H), 1.23 (t, 3H), 1.62 (m, 2H), 2.61 (t,
2H), 3.00-3.09 (m, 4H), 4.36 (t, 2H), 7.15 (d, 1H), 7.57 (t, 1H), 7.87 (d, 1H),
8.85 (s, 1H).

WO 94/29274

PCT/SE94/00552

-45-

Example VIII

Preparation of 3-propanoyl-4-chloro-8-(2-propylthiopropoxy)-quinoline

3-Propanoyl-4-chloro-8-(3-propylthiopropoxy) quinoline was synthesized
5 according to example V. Yield 5.5 g (87%).

¹H-NMR, 300 MHz, CDCl₃) 0.90 (t, 3H), 1.20 (t, 3H), 1.54 (m, 2H), 2.22
(m, 2H), 2.44 (t, 2H), 2.72 (t, 2H), 3.00 (q, 2H), 4.30 (t, 2H), 7.14 (d, 1H), 7.53
(t, 1H), 7.81 (d, 1H), 8.83 (s, 1H).

10

Example IX

Preparation of 3-butyryl-4-chloro-8-(3-methylthiopropoxy)-quinoline.

The title compound was synthesized according to the method in Example
15 V. Yield 2.9 g (91%) (hydrochloride).

¹H-NMR, 500 MHz, CDCl₃) 1.02 (t, 3H), 1.8 (m, 2H), 2.30 (s, 3H), 2.40 (m,
2H), 2.90 (t, 2H), 3.10 (t, 2H), 4.50 (t, 2H), 7.52 (d, 1H), 7.90-7.95 (m, 1H),
8.30-8.50 (m, 1H), 9.48 (s, 1H).

20

Example X

Preparation of 3-propanoyl-4-chloro-8-(3-methylthiopropoxy)quinoline.

The title compound was synthesized according to the method in Example
25 V. Yield 3.95 g (96%).

¹H-NMR, 300 MHz, CDCl₃) 1.28 (t, 3H), 2.1 (s, 3H), 2.3 (q, 2H), 2.8 (t, 2H),
3.08 (q, 2H), 4.38 (t, 2H), 7.23 (d, 1H), 7.6 (t, 1H), 7.9 (d, 1H), 8.9 (s, 1H).

WO 94/29274

PCT/SE94/00552

-46-

3. PREPARATION OF PHARMACEUTICAL FORMULATIONS

Pharmaceutical formulations containing a compound of the invention as active ingredient are illustrated in the following examples:

5

Formulation A. Syrup.

A syrup containing 1% (weight per volume) of active substance is prepared from the following ingredients:

10

Compound according to Example 2	1.0 g
Sugar, powder	30.0 g
Saccharine	0.6 g
Glycerol	5.0 g
15 Flavouring agent	0.05 g
Ethanol 96%	5.0 g
Distilled water q.s. to a final volume of	100 ml

20

Sugar and saccharine are dissolved in 60 g of warm water. After cooling the acid addition salt is dissolved in the sugar solution and glycerol and a solution of flavouring agents dissolved in ethanol are added. The mixture is diluted with water to a final volume of 100 ml.

25

The above given active substance may be replaced with other pharmaceutically acceptable acid addition salts.

Formulation B. Tablets

30

A tablet containing 50 mg of active compound is prepared from the following ingredients:

WO 94/29274

PCT/SE94/00552

-48-

Compound according to Example 2	4 g
Polyethylene glycol 400 for injection	400 g
Disodium hydrogen phosphate	q.s.
Sterile water to a final volume of	1.000 ml

5

Compound according to Example 2 is dissolved in polyethylene glycol 400 and 550 ml of water is added. pH of the solution is brought to pH 7.4 by adding a water solution of disodium hydrogen phosphate and water is added to a final volume of 1000 ml. The solution is filtered through a 0.22 μm filter and immediately dispensed into 10 ml sterile ampoules. The ampoules are sealed.

10

4. BIOLOGICAL TESTS

15 A. Inhibiting effect on acid secretion *in vitro* in isolated rabbit gastric glands was measured as described by Berglindh et al. (1976) Acta Physiol. Scand. 97, 401-414. The compounds according to Examples 1-12 had an IC_{50} value in the range of 0.5 to 6.0 μM . The compounds according to Examples 13-57 had an IC_{50} value in the range of 0.75 to 14 μM

20

B. Inhibiting effect on acid secretion *in vivo* in conscious female rats was measured according to the following method:

25 Female rats of the Sprague-Dawley strain were used. They were equipped with cannulated fistulae in the stomach (lumen) and the upper part of the duodenum, for collection of gastric secretions and administration of test substances, respectively. A fourteen days recovery period after surgery was allowed before testing commenced.

30 Before secretory tests, the animals were deprived of food but not water for 20 h. The stomach was repeatedly washed through the gastric cannula with tap water (37°C), and 6 ml of Ringer-Glucose given subcutaneously. Acid secretion was stimulated with infusion during 3 h (1.2 ml/h,

WO 94/29274

PCT/SE94/00552

—49—

subcutaneously) of pentagastrin and carbachol (20 and 110 nmol/kg h, respectively), during which time gastric secretions were collected in 30-min fractions. Test substances or vehicles were given intravenously or intraduodenally at 60 min after starting the stimulation, in a volume of 1.2 ml/h. Gastric juice samples were titrated to pH 7.0 with NaOH, 0.1 mole/L, and acid output calculated as the product of titrant volume and concentration.

Further calculations were based on group mean responses from 4-5 rats. The acid output during the periods after administration of test substances or vehicle were expressed as fractional responses, setting the acid output in the 30-min period preceding administration to 1.0. Percentage inhibition was calculated from the fractional responses elicited by test compound and vehicle. ED₅₀ values were obtained from graphical interpolation on log dose-response curves, or estimated from single-dose experiments assuming a similar slope for all dose-response curves.

The compounds 1-12 in Table 1 had an ED₅₀ value in the range of 1.0-12 µmole/kg. The results are based on gastric acid secretion during the second hour after drug/vehicle administration.


WO 94/29274

PCT/SE94/00552

-50-

CLAIMS

1. A compound of the general formula I



or a pharmaceutically acceptable salt of the said compound, wherein
R¹ is

- 20
- (a) C₁-C₆ alkyl,
 - (b) C₃-C₆ cykloalkyl, or
 - (c) C₃-C₆, C₁-C₆ cykloalkylalkyl;

R² is

- 25
- (a) H,
 - (b) C₁-C₆ alkyl,
 - (c) C₁-C₆ alkoxy, or
 - (d) halogen;

R³ is C₁-C₆ alkyl;

R⁴ is

- 30
- (a) H,
 - (b) C₁-C₄ alkyl,
 - (c) halogen, or
 - (d) OH;

m is an integer 2 or 3; and

WO 94/29274

PCT/SE94/00552

—51—

n is an integer 0, 1 or 2.

2. A compound according to claim 1, or a pharmaceutically acceptable salt of the said compound, wherein

5 R^1 is CH_3 , CH_2CH_3 , $CH_2CH_2CH_3$, $CH(CH_3)_2$, cyclopropyl or cyclopropylmethyl,

R^2 is CH_3 , CH_2CH_3 , $CH_2CH_2CH_3$, $CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, OCH_3 , OCH_2CH_3 , or halogen;

R^3 is CH_3 , CH_2CH_3 , $CH(CH_3)_2$ or $CH_2CH_2CH_3$; and

10 R^4 is H, CH_3 , CH_2CH_3 , halogen or OH.

3. A compound according to claim 2, or a pharmaceutically acceptable salt of the said compound, wherein

R^1 is CH_2CH_3 or $CH_2CH_2CH_3$;

15 R^2 is CH_3 , CH_2CH_3 , $CH(CH_3)_2$, OCH_3 , or Cl;

R^3 is CH_3 , CH_2CH_3 , or $CH_2CH_2CH_3$; and

R^4 is H, CH_3 , F, Cl or OH.

4. A compound according to claim 3, or a pharmaceutically acceptable salt of the said compound, which is one of the compounds listed in the Table below

25

30

No.	R^1	R^2	R^3	R^4	m	n
1	$CH_2CH_2CH_3$	CH_3	CH_3	H	2	0
2	$CH_2CH_2CH_3$	CH_3	CH_3	H	2	1
3	$CH_2CH_2CH_3$	CH_3	CH_3	H	2	2
4	$CH_2CH_2CH_3$	$CH(CH_3)_2$	CH_3	H	2	0
5	$CH_2CH_2CH_3$	$CH(CH_3)_2$	CH_3	H	2	1
6	$CH_2CH_2CH_3$	$CH(CH_3)_2$	CH_3	H	2	2
7	CH_2CH_3	CH_3	CH_3	H	2	0

WO 94/29274

PCT/SE94/00552

-52-

No.	R ¹	R ²	R ³	R ⁴	m	n
8	CH ₂ CH ₃	CH ₃	CH ₃	H	2	1
9	CH ₂ CH ₃	CH ₃	CH ₃	H	2	2
10	CH ₂ CH ₃	CH ₂ CH ₃	CH ₃	H	2	0
11	CH ₂ CH ₃	CH ₂ CH ₃	CH ₃	H	2	1
12	CH ₂ CH ₃	CH ₂ CH ₃	CH ₃	H	2	2
13	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	4-F	2	0
14	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	4-F	2	1
15	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	4-F	2	2
16	CH ₂ CH ₃	CH(CH ₃) ₂	CH ₃	H	2	0
17	CH ₂ CH ₂ CH ₃	CH ₂ CH ₃	CH ₃	H	2	0
18	CH ₂ CH ₂ CH ₃	CH ₂ CH ₃	CH ₃	H	2	1
19	CH ₂ CH ₂ CH ₃	CH ₂ CH ₃	CH ₃	H	2	2
20	CH ₂ CH ₃	CH ₃	CH ₂ CH ₂ CH ₃	H	2	0
21	CH ₂ CH ₃	CH ₃	CH ₂ CH ₂ CH ₃	H	2	1
22	CH ₂ CH ₃	CH ₃	CH ₂ CH ₂ CH ₃	H	2	2
23	CH ₂ CH ₃	CH ₃	CH ₂ CH ₂ CH ₃	H	3	0
24	CH ₂ CH ₃	CH ₃	CH ₂ CH ₂ CH ₃	H	3	1
25	CH ₂ CH ₃	CH ₃	CH ₂ CH ₂ CH ₃	H	3	2
26	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	4-OH	2	0
27	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	4-OH	2	1
28	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	4-OH	2	2
29	CH ₂ CH ₂ CH ₃	Cl	CH ₃	H	2	0
30	CH ₂ CH ₂ CH ₃	Cl	CH ₃	H	2	1
31	CH ₂ CH ₂ CH ₃	Cl	CH ₃	H	2	2
32	CH ₂ CH ₂ CH ₃	OCH ₃	CH ₃	H	2	0

WO 94/29274

PCT/SE94/00552

-53-

5

10

15

20

No.	R ¹	R ²	R ³	R ⁴	m	n
33	CH ₂ CH ₂ CH ₃	OCH ₃	CH ₃	H	2	1
34	CH ₂ CH ₂ CH ₃	OCH ₃	CH ₃	H	2	2
35	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	4-CH ₃	2	0
36	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	4-CH ₃	2	1
37	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	4-CH ₃	2	2
38	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	6-CH ₃	2	0
39	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	6-CH ₃	2	1
40	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	6-CH ₃	2	2
41	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	6-Cl	2	0
42	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	6-Cl	2	1
43	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	6-Cl	2	2
44	CH ₂ CH ₃	CH ₃	CH ₃	H	3	0
45	CH ₂ CH ₃	CH ₃	CH ₃	H	3	1
46	CH ₂ CH ₃	CH ₃	CH ₃	H	3	2
47	CH ₂ CH ₂ CH ₃	Cl	CH ₃	H	3	0
48	CH ₂ CH ₂ CH ₃	Cl	CH ₃	H	3	1
49	CH ₂ CH ₂ CH ₃	Cl	CH ₃	H	3	2
50	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	H	3	0
51	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	H	3	1
52	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	H	3	2
53	CH ₂ CH ₂ CH ₃	OCH ₃	CH ₃	H	3	0
54	CH ₂ CH ₂ CH ₃	OCH ₃	CH ₃	H	3	1
55	CH ₂ CH ₂ CH ₃	OCH ₃	CH ₃	H	3	2

25

5. The compound 3-butyryl-4-(2-methylphenylamino)-8-(2-methylsulfinylethoxy)quinoline or a pharmaceutical salt thereof.


WO 94/29274

PCT/SE94/00552


-54-

6. A process for the preparation of a compound as claimed in any one of claims 1-5, comprising

(a) reacting a compound of the general formula II



10 wherein R² and R⁴ is as defined in claim 1, with a compound of the general formula III



wherein R¹, R³, m and n are as defined in claim 1 and X is a leaving group, such as a halide, tosyloxy or mesyloxy; or

25 (b) preparing a compound of the formula I wherein R¹, R², R³, R⁴ and m are as defined in claim 1 and n is 1 or 2, by oxidizing a compound of formula I wherein R¹, R², R³, R⁴ and m are as defined in claim 1 and n is 0.

30 7. A process according to claim 6 wherein the said oxidation step is performed by (i) using an oxidizing agent, whereby oxidation takes place in a solvent such as halogenated hydrocarbons, alcohols, ethers or ketones; or (ii) carrying out the oxidation enzymatically by using an oxidizing

WO 94/29274

PCT/SE94/00552

—55—

enzyme; or (iii) carrying out the oxidation microbially by using a suitable microorganism.

5 8. A compound as claimed in any one of claims 1-5 for use in therapy.

9. A compound as claimed in any one of claims 1-5 for use in inhibition of gastric acid secretion and / or for treatment of gastrointestinal inflammatory diseases.

10 10. A pharmaceutical formulation containing a compound as claimed in any one of claims 1-5 as active ingredient.

11. Use of a compound as claimed in any one of claims 1-5 for the manufacture of a medicament for the inhibition of gastric acid secretion.

15 12. Use of a compound as claimed in any one of claims 1-5 for the manufacture of a medicament for the treatment of gastrointestinal inflammatory diseases.

20 13. Use of a compound as claimed in any one of claims 1-5 for the manufacture of a medicament for the treatment or prophylaxis of conditions involving infection by *Helicobacter pylori* of human gastric mucosa, wherein the said compound is adapted to be administered in combination with an antimicrobial agent.

25 14. A method for inhibiting gastric acid secretion which comprises administering to a mammal, including humans, in need of such inhibition an effective amount of a compound as claimed in any one of claims 1-5.

30 15. A method for the treatment of gastrointestinal inflammatory diseases which comprises administering to a mammal, including humans, in need of such treatment an effective amount of a compound as claimed in any one of claims 1-5.

WO 94/29274

PCT/SE94/00552

—56—


16. A method for the treatment or prophylaxis of conditions involving infection by *Helicobacter pylori* of human gastric mucosa, which comprises administering to a mammal, including humans, in need of such treatment an effective amount of a compound as claimed in any one of claims 1-5, wherein the said compound is administered in combination with an antimicrobial agent.

17. A pharmaceutical formulation for use in the inhibition of gastric acid secretion wherein the active ingredient is a compound according to any one of claims 1-5.

18. A pharmaceutical formulation for use in the treatment of gastrointestinal inflammatory diseases wherein the active ingredient is a compound according to any one of claims 1-5.

19. A pharmaceutical formulation for use in the treatment or prophylaxis of conditions involving infection by *Helicobacter pylori* of human gastric mucosa, wherein the active ingredient is a compound according to any one of claims 1-5 in combination with an antimicrobial agent.

20. A compound of the general formula III



III

WO 94/29274

PCT/SE94/00552

—57—

wherein R^1 represents C_1-C_6 alkyl; R^3 represents C_1-C_6 alkyl; m represents 2 or 3; n represents 0, 1 or 2; and X is a leaving group such as a halide, tosyloxy or mesyloxy.

1

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 94/00552

A. CLASSIFICATION OF SUBJECT MATTER

IPC 5: C07D 215/44, C07D 215/26, A61K 31/47

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 5: C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

REG, CA

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO, A1, 9212969 (SMITHKLINE BEECHAM INTERCREDIT B.V.), 6 August 1992 (06.08.92) ---	1-13, 17-20
X	EP, A1, 0330485 (SMITHKLINE BECKMAN INTERCREDIT B.V.), 30 August 1989 (30.08.89) ---	1-13, 17-20
X	EP, A2, 0416749 (SMITHKLINE BECKMAN INTERCREDIT B.V.), 13 March 1991 (13.03.91) ---	1-13, 17-20
X	EP, A1, 0259174 (SMITHKLINE & FRENCH LABORATORIES LIMITED), 9 March 1988 (09.03.88) ---	1-13, 17-20

☒ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"B" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

14 Sept 1994

Date of mailing of the international search report

21 -09- 1994

Name and mailing address of the ISA/
Swedish Patent Office

Box 5055, S-102 42 STOCKHOLM

Facsimile No. +46 8 666 02 86

Authorized officer

Irja Berlin

Telephone No. +46 8 782 25 00

2

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 94/00552

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US, A, 5143920 (ROBERT J. IFE ET AL), 1 Sept 1992 (01.09.92) --	1-13,17-20
X	US, A, 5082848 (ROBERT J. IFE ET AL), 21 January 1992 (21.01.92) --	1-13,17-20
X	WO, A1, 9312090 (SMITHKLINE BEECHAM INTERCREDIT B.V.), 24 June 1993 (24.06.93) -- -----	1-13,17-20

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 94/00552

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 14-16
because they relate to subject matter not required to be searched by this Authority, namely:
See PCT Rule 39.1(iv): Methods for treatment of the human or animal body by surgery or therapy, as well as diagnostic methods.
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐

The additional search fees were accompanied by the applicant's protest.

☐

No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

27/08/94

International application No.

PCT/SE 94/00552

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A1- 9212969	06/08/92	AP-A- 337 AU-B- 652028 AU-A- 1179992 CA-A- 2099117 CN-A- 1064268 EP-A- 0569396 JP-T- 6504541 NZ-A- 241408	08/04/94 11/08/94 27/08/92 30/07/92 09/09/92 18/11/93 26/05/94 26/05/94
EP-A1- 0330485	30/08/89	SE-T3- 0330485 AU-B- 606508 AU-A- 3181989 DE-T- 68906687 HU-B- 208814 IL-A- 89346 IL-A- 106072 JP-T- 2503318 PT-B- 89789 US-A- 5089504 WO-A- 8908104	07/02/91 22/09/89 02/12/93 28/01/94 25/01/94 12/04/94 11/10/90 31/05/94 18/02/92 08/09/89
EP-A2- 0416749	13/03/91	JP-A- 3086865 US-A- 5082841	11/04/91 21/01/92
EP-A1- 0259174	09/03/88	SE-T3- 0259174 AU-B- 598299 AU-A- 7874187 DE-A- 3777505 JP-T- 1500664 US-A- 4806549 US-A- 4806550 WO-A- 8801621	21/06/90 24/03/88 23/04/92 09/03/89 21/02/89 21/02/89 10/03/88
US-A- 5143920	01/09/92	AU-A- 3181789 DE-D,T- 68909107 EP-A,B- 0342775	22/09/89 17/02/94 23/11/89
US-A- 5082848	21/01/92	AU-A- 3011689 EP-A- 0331357	31/08/89 06/09/89
WO-A1- 9312090	24/06/93	AU-A- 3158593	19/07/93